arXiv:1203.2904v2 [math.CA] 6 Feb 2013

A density theorem in parameterized differential Galois theory.

Thomas Dreyfus

Abstract

We study parameterized linear differential equations with coefficients depending meromorphically upon the parameters. As a main result, analogously to the unparameterized density theorem of Ramis, we show that the parameterized monodromy, the parameterized exponential torus and the parameterized Stokes operators are topological generators in Kolchin topology, for the parameterized differential Galois group introduced in [CS]. We prove an analogous result for the global parameterized differential Galois group, which generalizes a result in [MS11]. In [MS11] the authors give also a necessary condition on a group for being a global parameterized differential Galois group: as a corollary of the density theorem, we prove that their condition is also sufficient. As an application, we give a characterization of completely integrable equations, and we give a partial answer to a question of Sibuya about the transcendence properties of a given Stokes matrix. Moreover, using a parameterized Turrittin's theorem, we show that the Galois group descents to a smaller field, whose field of constants is not differentially closed.

Introduction

Let us consider a linear differential system of the form

$$\partial_X Y(X) = A(X)Y(X),$$

where $\partial_X = \frac{d}{dX}$, and $A(X)$ is an $m \times m$ matrix whose entries are germs of meromorphic functions in a neighborhood of a point, say 0 to fix ideas. The differential Galois group, which measures the algebraic dependencies among the solutions, can be viewed as an algebraic subgroup of $\mathrm{GL}_m(\mathbb{C})$ via the injective group morphism

$$\begin{array}{cccc} \rho_U : & Gal & \longrightarrow & \mathrm{GL}_m(\mathbb{C}) \\ & \sigma & \mapsto & U(X)^{-1}\sigma(U(X)), \end{array}$$

where $U(X)$ is some fundamental solution (i.e: an invertible matrix solution).

Let $U(X)$ be a fundamental solution of $\partial_X Y(X) = A(X)Y(X)$. The linear differential equation is said to be regular singular at 0 if there exists an invertible matrix $P(X)$ whose entries are germs of meromorphic functions, such that $Z(X) = P(X)U(X)$ satisfies

$$\partial_X Z(X) = \frac{A_0}{X} Z(X),$$

where A_0 is a matrix with complex entries. In this case, $Z(X)$ involves multivalued functions.

2010 Mathematics Subject Classification Primary 34M03, 12H05, 34M30, 34M15, 34M40, 34M50

Work partially supported by ANR, contract ANR-06-JCJC-0028

Analytic continuation of $Z(X)$ along any path γ sufficiently close to 0 and centered at 0 yields another fundamental solution $Z(X)M_{\gamma}$, and the matrix M_{γ} with complex entries, which is a monodromy matrix, does not depend on the choice of γ. The Schlesinger theorem says that the Zariski closure of the group generated by the monodromy matrix is the Galois group. In the general case (i.e: in presence of irregular singularities), the monodromy is no longer sufficient to provide a complete collection of topological generators. Ramis has shown that the monodromy, the exponential torus and the Stokes operators are topological generators for the Galois group in the Zariski topology.

More recently, a Galois theory for parameterized linear differential equations of the form

$$\partial_X Y(X,t) = A(X,t)Y(X,t),$$

where $t=(t_1,\dots,t_n)$ are parameters, has been developed in [CS]. See also [HS, L]. Namely, the Galois group, which measures the $(\partial_{t_1},\dots,\partial_{t_n})$-differential and algebraic dependencies among the solutions, can be seen as a differential group in the sense of Kolchin, that is a group of matrices whose entries lie in a differential field and satisfy a set of polynomial differential equations in the variables $t_1,\dots,t_n$. To be applied, the theory from [CS] requires the field of constants with respect to ∂_X to be of characteristic 0 and differentially closed (see §2.1). The drawback of this latter assumption is that a differentially closed field is a very big field, and cannot be interpreted as a field of functions. The main result of this paper is a parameterized analogue of the density theorem of Ramis: we give topological generators for the Galois group, for the Kolchin topology (in which closed sets are zero sets of differential algebraic polynomials).

There is a link between the parameterized differential Galois theory and isomonodromy for equations with only regular singular poles (see [CS, MS, MS11]). Let $\mathcal{D}(t,r)$ be an open polydisc in $\mathbb{C}^n$, let $\mathcal{D}$ be an open subset of $\mathbb{C}$, and let $A(X,t)$ be a matrix whose entries are analytic on $\mathcal{D}\times\mathcal{D}(t,r)$. We consider open disks D_j that cover $\mathcal{D}$, and a solution $U_j(X,t)$ of $\partial_X Y(X,t) = A(X,t)Y(X,t)$ analytic on $D_j\times\mathcal{D}(t,r)$. If $D_i\cap D_j\neq\varnothing$, we define $C_{i,j}(t)=U_i(X,t)^{-1}U_j(X,t)$, the connection matrices. The parameterized linear differential equation $\partial_X Y(X,t)=A(X,t)Y(X,t)$ is said to be isomonodromic if, there is a choice of D_i covering $\mathcal{D}$, and of the solutions $U_i(X,t)$ of $\partial_X Y(X,t)=A(X,t)Y(X,t)$ analytic on $D_i\times\mathcal{D}(t,r)$, such that the connection matrices are independent of t. In this case, the matrix of the monodromy is constant on the polydisc $\mathcal{D}(t,r)$. In the regular singular case, the following statements are equivalent (see [CS], Propositions 5.3 and 5.4).

- The Galois group is conjugate, over a differentially closed field (see Definition 2.2), to a group of constant matrices.
- The parameterized linear differential equation is isomonodromic.
- The parameterized linear differential equation is completely integrable (see Definition 3.1).

In [GO], the authors study completely integrable parameterized linear differential equations using differential Tannakian categories. In particular, they prove that the notion of integrability with respect to all the parameters is equivalent to the notion of integrability with respect to each parameter separately, which generalizes [D], Proposition 9. Furthermore, they improve Proposition 3.9 in [CS] by avoiding the assumption that the field of constants is differentially closed.

As an application of our main result, the parameterized density theorem, we improve Proposition 3.9 in [CS] (see Remark 3.5): the parameterized linear differential equation is completely integrable if and only if the topological generators for the Galois group given in the parameterized density theorem are conjugate to constant matrices over a field of meromorphic functions. Notice that this latter is not differentially closed.

The article is organized as follows. In the first section we study parameterized linear differential systems from an analytic point of view. The parameters will vary in U, a non-empty polydisc of $\mathbb{C}^n$. Let $t=(t_1,\dots,t_n)\in U$ denote the multiparameter. Let $\mathcal{M}_U$ be the field of meromorphic functions on U and let $\hat{K}_U=\mathcal{M}_U[[X]][X^{-1}]$. The Hukuhara-Turrittin's theorem in this case gives the following result:

Proposition (see Proposition 1.1 below). *Let* $\partial_X Y(X,t)=A(X,t)Y(X,t)$*, with* $A(X,t)\in \mathrm{M}_m(\hat{K}_U)$ *(that is a* $m\times m$ *matrix with entries in* $\hat{K}_U$*). Then, there exists a polydisc* $\varnothing\neq U'\subset U$*,* $\nu\in\mathbb{N}^*$*, such that we have a fundamental solution* $F(X,t)$ *of the form:*

$$F(X,t)=\hat{H}(X,t)X^{L(t)}e^{Q(X,t)},$$

where:

- $\hat{H}(X,t)\in \mathrm{GL}_m(\hat{K}_{U'}[X^{1/\nu}])$.
- $L(t)\in \mathrm{M}_m(\mathcal{M}_{U'})$.
- $e^{Q(X,t)}=\mathrm{Diag}(e^{q_i(X,t)})$*, with* $q_i(X,t)\in X^{-1/\nu}\mathcal{M}_{U'}[X^{-1/\nu}]$.
- *Moreover, we have* $X^{L(t)}e^{Q(X,t)}=e^{Q(X,t)}X^{L(t)}$.

Combined with a result of Schaëfke (see Remark 1.3), we have a sufficient condition on $t_0\in U$, to have a fundamental solution in the same form as above with $t_0\in U'$.

In §1.3, we briefly review the Stokes phenomenon in the unparameterized case. We have solutions, which are analytic in some sector and Gevrey asymptotic to the formal part of the solution in the Hukuhara-Turrittin's canonical form. The fact that various asymptotic solutions do not glue to a single solution on the Riemann surface of the logarithm is called the Stokes phenomenon.

Let $\varnothing\neq U$ be a polydisc of $\mathbb{C}^n$ and let $f(X,t)=\sum f_i(t)X^i\in\hat{K}_U$. We say that $f(X,t)\in\mathcal{O}_U$ if for all $t\in U$, $X\mapsto\sum f_i(t)X^i$ is a germ of meromorphic function. Notice that if $f(X,t)\in\mathcal{O}_U\subset\mathcal{M}_U[[X]][X^{-1}]=\hat{K}_U$, then the X-coefficients of $f(X,t)$ are analytic on U.

In §1.4, we study the Stokes phenomenon of equations of the form $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$. In particular, we prove that the asymptotic solutions depend analytically upon the parameters.

In the second section, we use the parameterized Hukuhara-Turrittin's theorem to deduce some Galois theoretic properties of the systems. We first recall some facts from [CS] about parameterized differential Galois theory. The problem is that the theory in [CS] cannot be applied here, since $\mathcal{M}_U$, our field of constants with respect to ∂_X, is a field of functions that are meromorphic in $t_1,\dots,t_n$, and this field is not differentially closed (see §2.1). In the papers [GGO, W], the authors prove the existence of parameterized Picard-Vessiot extensions under weaker assumptions than in [CS]. We do not use these latter results because we need a parameterized Turrittin's theorem, which proves directly the descent here, in order to study the parameterized Stokes phenomenon. In §2.4, we consider $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$. We state and show the main result:

Parameterized analogue of the density theorem of Ramis (Theorem 2.17). *The group generated by the parameterized monodromy, the parameterized exponential torus and the parameterized Stokes operators is dense in the parameterized Galois group for the Kolchin topology.*

Then, we turn to the global case. We consider equations with coefficients in $\mathcal{M}_U(X)$ and study their global Galois group. We prove a density theorem in this global setting, see Theorem 2.21. The proof in the unparameterized case can be found in [M]. In §2.6, we give various examples of computation.

In the third section, we give three applications. First, we prove a criterion for the integrability of differential systems (see Definition 3.1):

Proposition (see Proposition 3.3 below). *Let* $A(X,t) \in \mathrm{M}_m(\mathcal{M}_U(X))$. *The linear differential equation* $\partial_X Y(X,t) = A(X,t)Y(X,t)$ *is completely integrable if and only if there exists a fundamental solution such that the matrices of the parameterized monodromy, the parameterized exponential torus and the parameterized Stokes operators for all the singularities are constant.*

As a second application, we give a partial answer to a question of Sibuya (see [Si]), regarding the differential transcendence properties of a Stokes matrix of the parameterized linear differential equation:

$$\begin{pmatrix} \partial_X Y(X,t) \\ \partial_X^2 Y(X,t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ X^3+t & 0 \end{pmatrix} \begin{pmatrix} Y(X,t) \\ \partial_X Y(X,t) \end{pmatrix}.$$

Sibuya was asking whether an entry of a given Stokes matrix at infinity is ∂_t-differentially transcendental (i.e: satisfies no differential polynomial equation). We prove that it is at least not ∂_t-finite (i.e: that it satisfies no linear differential equations).

As a last application, we deal with the inverse problem. We prove that if G is the global parameterized differential Galois group of some equation having coefficients in $k(X)$ (see §3.3), then G contains a finitely generated Kolchin dense subgroup. The converse of this latter assertion has been proved in Corollary 5.2 in [MS11], and we obtain a result on the inverse problem:

Theorem (see Theorem 3.12 below). *G is the global parameterized differential Galois group of some equation having coefficients in $k(X)$ if and only if G contains a finitely generated Kolchin-dense subgroup.*

In the appendix, we prove the following result.

Theorem (see Theorem A.1 below). *Let* $\partial_X Y(X,t) = A(X,t)Y(X,t)$, *with* $A(X,t) \in \mathrm{M}_m(\hat{K}_U)$. *Then, there exists a polydisc* $\varnothing \neq U' \subset U$, *such that we have a fundamental solution* $F(X,t)$ *of the form:*

$$F(X,t) = \hat{P}(X,t)X^{C(t)}e^{Q(X,t)},$$

where:

- $\hat{P}(X,t) \in \mathrm{GL}_m(\hat{K}_{U'})$.
- $C(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$.
- $e^{Q(X,t)} = \mathrm{Diag}(e^{q_i(X,t)})$, *with* $q_i(X,t) \in X^{-1/\nu}\mathcal{M}_{U'}[X^{-1/\nu}]$, *for some* $\nu \in \mathbb{N}^*$.

This theorem is not necessary for the proof of the main result of the paper, this is the reason why we give the proof in the appendix. Notice that contrary to Proposition 1.1, the entries of the formal part are not ramified. On the other hand, $X^{C(t)}$ and $e^{Q(X,t)}$ do not commute anymore.

Acknowledgments. This paper was prepared during my thesis, supported by the region Ile de France. I want to thank my advisor, Lucia Di Vizio for her helpful comments and interesting discussions we had during the preparation of this paper. I also want to thank the organizers of seminars in Toulouse, Strasbourg, Lille and Grenoble, and of the meeting "Introduction aux groupes quantiques (ANR-qdiff)" in Marseille, for making it possible for me to present the results contained in this paper. I want to thank Jean-Pierre Ramis, Guy Casale, Reinhard Schaëfke, Daniel Bertrand and Michael F. Singer for pointing some mistakes and inaccuracies in this paper. Michael F. Singer in particular suggested the contribution of this paper to Theorem 3.12. I certainly thank Carlos E. Arreche and Claude Mitschi for the read-through. Lastly, I heartily thank the two anonymous referees who spent a great lot of time and effort to help me make the present paper readable.

1. Local analytic linear differential system depending upon parameters.

In §1.1, we define the field to which the entries of the fundamental solution, in the Hukuhara-Turrittin's canonical form, will belong. In §1.2, we prove a parameterized version of the Hukuhara-Turrittin's theorem. In §1.3, we briefly review the Stokes phenomenon in the unparameterized case. In §1.4, we study the Stokes phenomenon in the parameterized case.

1.1 Definition of the fields.

Let us consider a linear differential system of the form $\partial_X Y(X) = A(X)Y(X)$, where $A(X)$ is a $m \times m$ matrix whose entries belongs to $\mathbb{C}[[X]][X^{-1}]$. We know we can find a formal fundamental solution in the Hukuhara-Turrittin's canonical form $\hat{H}(X)X^L e^{Q(X)}$ (see Theorem 3.1 in [VdPS]), where:

- $\hat{H}(X)$ is a matrix of formal power series in $X^{1/\nu}$ for some $\nu \in \mathbb{N}^*$.
- $L \in \mathrm{M}_m(\mathbb{C})$.
- $Q(X) = \mathrm{Diag}(q_i(X))$, with $q_i(X) \in X^{-1/\nu}\mathbb{C}[X^{-1/\nu}]$.
- Moreover, we have $X^L e^{Q(X)} = e^{Q(X)} X^L$.

Let $\varnothing \neq U$ be a polydisc of $\mathbb{C}^n$ and let $\hat{K}_U$ defined in page 3. We want to construct a field containing a fundamental set of solutions of

$$\partial_X Y(X,t) = A(X,t)Y(X,t),$$

with $A(X,t) \in \mathrm{M}_m(\hat{K}_U)$. Let $\Delta_t = \{\partial_{t_1}, \dots, \partial_{t_n}\}$ and let

$$\mathbf{E}_U = \bigcup_{\nu \in \mathbb{Q}^{>0}} X^{\frac{-1}{\nu}} \mathcal{M}_U [X^{\frac{-1}{\nu}}].$$

We define formally the (∂_X, Δ_t)-ring

$$\hat{K}_U \left[\log, \left(X^{a(t)} \right)_{a(t) \in \mathcal{M}_U}, \left(e(q(X,t)) \right)_{q(X,t) \in \mathbf{E}_U} \right],$$

with the following rules:

(i) The symbols log, $\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}$ and $(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}$ only satisfy the following relations:

$$\begin{array}{rclrcl} X^{a(t)+b(t)} & = & X^{a(t)}X^{b(t)}, & e(q_1(X,t)+q_2(X,t)) & = & e(q_1(X,t))e(q_2(X,t)),\\ X^{a} & = & X^{a}\in\hat{K}_U \text{ for } a\in\mathbb{Z}, & e(0) & = & 1. \end{array}$$

(ii) The differentiation in the (∂_X,Δ_t)-ring is given by

$$\begin{array}{rclrclrcl} \partial_X \log & = & X^{-1}, & \partial_X X^{a(t)} & = & \frac{a(t)}{X}X^{a(t)}, & \partial_X e(q(X,t)) & = & \partial_X(q(X,t))e(q(X,t)),\\ \partial_{t_i}\log & = & 0, & \partial_{t_i}X^{a(t)} & = & \partial_{t_i}(a(t))\log X^{a(t)}, & \partial_{t_i}e(q(X,t)) & = & \partial_{t_i}(q(X,t))e(q(X,t)). \end{array}$$

The intuitive interpretation of these symbols are $\log=\log(X)$, $X^{a(t)}=e^{a(t)\log(X)}$ and $e(q(X,t))=e^{q(X,t)}$. Let $f(X,t)$ be one these latter functions. Then $f(X,t)$ has a natural interpretation as an analytic function on $\widetilde{\mathbb{C}}\times U$, where $\widetilde{\mathbb{C}}$ is the Riemann surface of the logarithm. We will use the analytic function instead of the symbol when we will consider asymptotic solutions (see §1.3 and §1.4). For the time being, however, we see them only as symbols.

We can see (Proposition 3.22 in[VdPS]) that the (∂_X,Δ_t)-ring

$$\hat{K}_U[\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U},(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}]$$

is contained in an integral domain. Therefore, we may consider the (∂_X,Δ_t)-fields:

$$K_{F,U}=\mathcal{M}_U\left(\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}\right),$$

$$\hat{K}_{F,U}=\hat{K}_U\left(\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}\right),$$

and

$$\widehat{\mathbf{K}_U}=\hat{K}_U\left(\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}\right).$$

In the definition of the fields $K_{F,U}$ and $\hat{K}_{F,U}$, the subscript F stands for Fuchsian. We have defined (∂_X,Δ_t)-fields where all the derivations commute with each other. We have the following inclusions of (∂_X,Δ_t)-fields:

$$\begin{tikzcd} & K_{F,U} \arrow[dr] & & \\ \mathcal{M}_U \arrow[ur] \arrow[r] & \hat{K}_U \arrow[r] & \hat{K}_{F,U} \arrow[r] & \widehat{\mathbf{K}_U}. \end{tikzcd}$$

1.2 Hukuhara-Turrittin's theorem in the parameterized case.

The goal of this subsection is to give the parameterized version of the Hukuhara-Turrittin's theorem. We prove a slightly stronger result, which is not needed in the paper in the appendix, see Theorem A.1.

Proposition 1.1. *Let $\varnothing\neq U$ be a polydisc of $\mathbb{C}^n$ and consider*

$$\partial_X Y(X,t)=A(X,t)Y(X,t),$$

with $A(X,t)\in\mathrm{M}_m(\hat{K}_U)$. There exists a polydisc $\varnothing\neq U'\subset U$, such that we have a fundamental solution $F(X,t)\in\mathrm{GL}_m(\widehat{\boldsymbol{K}_{U'}})$ of the form

$$F(X,t)=\hat{H}(X,t)X^{L(t)}e(Q(X,t)),$$

where:

- *$\hat{H}(X,t) \in \mathrm{GL}_m(\hat{K}_{U'}[X^{1/\nu}])$, for some $\nu \in \mathbb{N}$.*
- *$L(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$ and $X^{L(t)} \in \mathrm{GL}_m(K_{F,U'})$ satisfy*

$$\partial_X X^{L(t)} = \frac{L(t)}{X} X^{L(t)} = X^{L(t)} \frac{L(t)}{X}.$$

- *$e(Q(X,t)) = \mathrm{Diag}(e(q_i(X,t)))$, with $q_i(X,t) \in \boldsymbol{E}_{U'}$.*
- *Moreover, we have $e(Q(X,t))X^{L(t)} = X^{L(t)}e(Q(X,t))$.*

Furthermore, if $A(X,t) \in \mathrm{M}_m(\mathcal{O}_U)$, there exists a polydisc $\varnothing \neq U'' \subset U'$, such that the entries of the X-coefficients of $\hat{H}(X,t)$ are all analytic on U''.

Example 1.2 ([Sch], Introduction). If we consider

$$X^2 \partial_X Y(X,t) = \begin{pmatrix} t & 1 \\ X & 0 \end{pmatrix} Y(X,t),$$

we get the solution

$$\left(\begin{pmatrix} 1 & 1 \\ 0 & -t \end{pmatrix} + O(X) \right) \begin{pmatrix} X^{\frac{1}{t}} e^{\frac{-t}{X}} & 0 \\ 0 & X^{\frac{-1}{t}} \end{pmatrix}, \tag{1}$$

for $t \neq 0$ and the solution

$$\begin{pmatrix} 1 & 1 \\ X^{1/2} & -X^{1/2} \end{pmatrix} \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + O(X^{1/2}) \right) \begin{pmatrix} X^{\frac{1}{4}} e^{-X^{-1/2}} & 0 \\ 0 & X^{\frac{1}{4}} e^{X^{-1/2}} \end{pmatrix},$$

for $t = 0$. The latter is not the specialization of (1) at $t = 0$. The problem is that the level of the unparameterized system (see §1.3 for the definition) at $t = 0$ is 1 and the level of the unparameterized system for $t \neq 0$ is $\frac{1}{2}$. This example shows that we cannot get a solution in the parameterized Hukuhara-Turrittin's form, that remains valid for all values of the parameter t. This is the reason why we have to remove a subset of the parameter-space.

Remark 1.3. Similar results to Proposition 1.1 have been proved in Theorem 10.1 of [BV] and Theorem 4.2 of [Sch]. For all t, let $\hat{H}_t(X)X^{L_t}e(Q(X,t))$ be a solution given in the classical Hukuhara-Turrittin's canonical form. In order to make precise the dependence on t, the author of [Sch] makes three additional assumptions; namely, that there exists $U \subset \mathbb{C}^n$ an open connected subset such that:

- The $q_i(X,t)$ are analytic functions in $t \in U$.
- The degree in X^{-1} of $q_i(X,t) - q_j(X,t)$ is independent of t in U.
- For all $t \in U$, $q_j(X,t) \neq q_i(X,t)$, if $i \neq j$.

Under these assumptions, Schaëfke concludes that, for all $t_0 \in U$, there exists an open neighborhood $\varnothing \neq U' \subset U$ of t_0 in the t-plane, such that there exists a solution $\hat{H}(X,t)X^{L(t)}e(Q(X,t)) \in \mathrm{GL}_{\mathrm{m}}(\widehat{\mathbf{K}_{U'}})$ with the same properties as in Proposition 1.1. Notice that Schaëfke gives a computationable necessary and sufficient condition for well behaved exponential part, see Theorem 5.2 in [Sch].

Remark 1.4. Algebraic functions over $\mathcal{M}_U$ can be seen as elements of $\mathcal{M}_{U'}$, for some non-empty $U' \subset U$. Therefore, a finite extension of $\mathcal{M}_U$ can be embedded in $\mathcal{M}_{U'}$ for a convenient choice of $U' \subset U$. We will use this fact in the rest of the paper.

Before proving Proposition 1.1, we prove the lemma:

Lemma 1.5. *Let $\varnothing \neq U$ be a polydisc of $\mathbb{C}^n$ and let $L(t) \in \mathrm{M}_m(\mathcal{M}_U)$. There exists a polydisc $\varnothing \neq U' \subset U$, and $X^{L(t)} \in \mathrm{GL}_m(K_{F,U'})$ satisfying*

$$\partial_X X^{L(t)} = \frac{L(t)}{X} X^{L(t)} = X^{L(t)} \frac{L(t)}{X}.$$

Proof. Consider $L(t)$ as a matrix with entries in $\overline{\mathcal{M}_U}$, the algebraic closure of $\mathcal{M}_U$. There exists a polydisc $\varnothing \neq U' \subset U$, such that the size of the Jordan block of $L(t)$ does not depend on t on U' and such that the eigenvalues of $L(t)$ belongs to $\mathcal{M}_{U'}$. Let $L(t) = P(t)(D(t)+N(t))P^{-1}(t)$, with $D(t) = \mathrm{Diag}(d_i(t))$, $d_i(t) \in \mathcal{M}_U$, $N(t)$ nilpotent, $D(t)N(t) = N(t)D(t)$ and $P(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$ be the Jordan decomposition of $L(t)$.

Then, the matrix $X^{L(t)} = P(t)\mathrm{Diag}(X^{d_i(t)})e^{N(t)\log}P^{-1}(t)$ belongs to $\mathrm{GL}_m(K_{F,U'})$ and $X^{L(t)}$ satisfies

$$\partial_X X^{L(t)} = \frac{L(t)}{X} X^{L(t)} = X^{L(t)} \frac{L(t)}{X}.$$

□

Proof of Proposition 1.1. The Turrittin's theorem (see Theorem 3.1 in [VdPS]) is valid for any linear differential system with entries in $K = C[[X]][X^{-1}]$, where C is any algebraically closed fields of characteristic 0 such that C is the field of constants of K with respect to ∂_X (i.e: $C = \{a \in K | \partial_X(a) = 0\}$) and $\partial_X X = 1$.

Let us consider $L(t) \in \mathrm{M}_m(\overline{\mathcal{M}_U})$ and $Q(X,t) = \mathrm{Diag}(q_i(X,t))$, with $q_i(X,t) \in X^{-1/\nu}\overline{\mathcal{M}_U}[X^{-1/\nu}]$ for some $\nu \in \mathbb{N}$. Because of Remark 1.4, there exists a polydisc $\varnothing \neq U' \subset U$, such that $L(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$ and $q_i(X,t) \in \mathbf{E}_{U'}$. Hence, there exists a polydisc $\varnothing \neq U' \subset U$ such that the Turrittin's theorem gives the fundamental solution

$$F(X,t) = \hat{H}(X,t)X^{L(t)}e(Q(X,t)),$$

where:

- $\hat{H}(X,t) \in \mathrm{GL}_m(\overline{\mathcal{M}_{U'}}[[X^{1/\nu}]][X^{-1/\nu}])$, for some $\nu \in \mathbb{N}$.
- $L(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$ and $X^{L(t)} \in \mathrm{GL}_m(K_{F,U'})$ satisfy
$$\partial_X X^{L(t)} = \frac{L(t)}{X} X^{L(t)} = X^{L(t)} \frac{L(t)}{X}.$$
- $e(Q(X,t)) = \mathrm{Diag}(e(q_i(X,t)))$, with $q_i(X,t) \in \mathbf{E}_{U'}$.
- Moreover, we have $e(Q(X,t))X^{L(t)} = X^{L(t)}e(Q(X,t))$.

Let us prove now that $\hat{H}(X,t) \in \mathrm{GL}_m(\hat{K}_{U'}[X^{1/\nu}])$. The matrix

$$F(X,t) = \hat{H}(X,t)X^{L(t)}e(Q(X,t))$$

satisfies the parameterized linear differential equation

$$\partial_X F(X,t) = A(X,t)F(X,t),$$

and the matrix $X^{L(t)}e(Q(X,t))$ satisfies parameterized linear differential equation:

$$\partial_X X^{L(t)}e(Q(X,t)) = \left(L(t) + \partial_X e(Q(X,t))\right) X^{L(t)}e(Q(X,t)) = X^{L(t)}e(Q(X,t)) \left(L(t) + \partial_X e(Q(X,t))\right).$$

Hence,

$$\partial_X \hat{H}(X,t) = A(X,t)\hat{H}(X,t) - \hat{H}(X,t)\left(L(t) + \partial_X e(Q(X,t))\right).$$

We write $\hat{H}(X,t)$ as a column vector $\widetilde{H}(X,t)$ of size m^2. Let $C(X,t)\in \mathrm{M}_{m^2}(\hat{K}_{U'}[X^{1/\nu}])$, with $\nu\in\mathbb{N}^*$, such that $\widetilde{H}(X,t)$ satisfies the parameterized linear differential system:

$$\partial_X \widetilde{H}(X,t)=C(X,t)\widetilde{H}(X,t).$$

Let us write $\widetilde{H}(X,t)=\sum\limits_{i\geqslant N}\widetilde{H}_i(t)X^{i/\nu}$ and $C(X,t)=\sum\limits_{i\geqslant M}C_i(t)X^{i/\nu}$, where $M,N\in\mathbb{N}$. Then, by identifying the coefficients of the $X^{i/\nu}$-terms of the power series in the equation $\partial_X\widetilde{H}(X,t)=C(X,t)\widetilde{H}(X,t)$, we find that:

$$((i+1)\mathrm{Id}-C_{-1}(t))\widetilde{H}_{i+1}(t)=\sum_{l=N,l\neq i+1}^{i-M}C_{i-l}(t)\widetilde{H}_l(t).$$

The fact that $\hat{H}(X,t)\in \mathrm{GL}_m(\hat{K}_{U'}[X^{1/\nu}])$ is now clear.

Assume now that $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$. Let $\varnothing\neq U''\subset U'$ such that for $0\neq X$ fixed, the entries of the X-coefficients of $L(t)+\partial_X e(Q(X,t))$ are analytic on U''. Then, the entries of the X-coefficients of $C(X,t)$ are all analytic on U''. Hence, the entries of the X-coefficients of $\hat{H}(X,t)$ are all analytic on U''. □

Remark 1.6. If we take a smaller polydisc $\varnothing\neq U$, we may assume assume that if we consider $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$, then the fundamental solution of Proposition 1.1 belongs to $\mathrm{GL}_m(\widehat{\mathbf{K}_U})$, and the entries of the X-coefficients of $\hat{H}(X,t)$ are all analytic on U.

1.3 Review of the Stokes phenomenon in the unparameterized case.

In this subsection we will briefly review the Stokes phenomenon in the unparameterized case. See [M95, LR90, LR95, CR, S09] or Chapter 8 of [VdPS] for more details. We will generalize some results concerning the summation of divergent series in the parameterized case in §1.4. First we treat the example of the Euler equation:

$$X^2\partial_X Y(X)+Y(X)=X,$$

which admits as a solution the formal series: $\hat{f}(X)=\sum_{n\geqslant 0}(-1)^n n! X^{n+1}$. Classical methods of differential equations give another solution:

$$f(X)=\int_0^X e^{1/X}e^{-1/t}\frac{dt}{t}=\int_0^\infty \frac{1}{1+u}e^{-u/X}du,$$

where $1/t-1/X=u/X$. The solution $\hat{f}(X)$ is divergent and the solution $f(X)$ can be extended to an analytic function on the sector:

$$V=\left\{X\in\widetilde{\mathbb{C}}\,\middle|\,\arg(X)\in\left]\frac{-3\pi}{2},\frac{+3\pi}{2}\right[\right\}.$$

On this sector, $f(X)$ is 1-Gevrey asymptotic to $\hat{f}(X)$: for every closed subsector W of V, there exists $A_W\in\mathbb{R}$, $\varepsilon>0$ such that for all N and all $X\in W$ with $|X|<\varepsilon$,

$$\left|f(X)-\sum_{n\geqslant 0}^{N-1}(-1)^n n! X^{n+1}\right|\leqslant (A_W)^{N+1}(N+1)!|X|^{N+1}.$$

We can also consider $f(e^{2i\pi}X)$, which is an asymptotic solution on the sector:

$$V'=\left\{X\in\widetilde{\mathbb{C}}\,\middle|\,\arg(X)\in\left]\frac{\pi}{2},\frac{7\pi}{2}\right[\right\}.$$

The two asymptotic solutions do not glue to a single asymptotic solution on $V \cup V'$. In fact, the residue theorem implies that the difference in $V \cap V'$ of the two asymptotic solutions is:

$$2i\pi e^{1/X}.$$

The fact that various asymptotic solutions do not glue to a single analytic solution is called Stokes phenomenon.

More generally, let us consider a linear differential equation $\partial_X Y(X) = A(X)Y(X)$, such that the entries of $A(X)$ are germs of meromorphic functions in a neighborhood of 0. Let $\hat{H}(X)X^L e(Q(X))$, with $Q(X) = \mathrm{Diag}(q_i(X))$, be a fundamental solution in Hukuhara-Turrittin's canonical form. Since for all $k \in \mathbb{N}$, $\hat{H}(X)X^L e(Q(X)) = \hat{H}(X)\mathrm{Diag}(X^k)X^{L-k\mathrm{Id}}e(Q(X))$, we may assume that $\hat{H}(X)$ has no pole at $X = 0$. The levels of $\partial_X Y(X) = A(X)Y(X)$ are the degrees in X^{-1} of the $q_i(X) - q_j(X)$ (the levels are positive rational numbers). Consider $q(X) = q_k X^{-k/\nu} + \cdots + q_1 X^{-1/\nu} \in X^{-1/\nu}\mathbb{C}[X^{-1/\nu}]$ with $\nu \in \mathbb{N}$. The real number d is called singular for $q(X)$ if $q_k e^{-idk/\nu}$ is a positive real number. This correspond to the arguments d such that $r \mapsto e^{q(re^{id})}$ increases fastest as r tends to 0^+. The singular directions of $\partial_X Y(X) = A(X)Y(X)$ (we will write singular directions when no confusion is likely to arise) are the real number that are singular for one of the $q_i(X) - q_j(X)$, with $i \neq j$. Notice that the set of singular directions is finite modulo $2\pi\nu$ for some $\nu \in \mathbb{N}$. Let $k_1 < \cdots < k_r$ be the levels of the linear differential equation. There exists a decomposition $\hat{H}(X) = \hat{H}_{k_1}(X) + \cdots + \hat{H}_{k_r}(X)$, such that for d not a singular direction, there exists an unique r-tuple of matrices $(H^d_{k_1}(X), \dots, H^d_{k_r}(X))$, such that $H^d_{k_i}(X)$ is analytic on the sector

$$V_d = \left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left] d - \frac{\pi}{2k_i}, d + \frac{\pi}{2k_i} \right[\right\},$$

and is k_i-Gevrey asymptotic to $\hat{H}_{k_i}(X) = \sum_{n \in \mathbb{N}} \hat{H}_{n,k_i} X^n$ on V_d: for every closed subsector W of V_d, there exists $A_W \in \mathbb{R}$, $\varepsilon > 0$ such that for all N and all $X \in W$ with $|X| < \varepsilon$,

$$\left| H^d_{k_i}(X) - \sum_{n \geqslant 0}^{N-1} \hat{H}_{n,k_i} X^n \right| \leqslant (A_W)^N \Gamma\left(1 + \frac{N}{k_i}\right) |X|^N,$$

where Γ denotes the Gamma function. Until the end of the paper, we will denote a fixed determination of the complex logarithm by $\log(X)$. Furthermore, the matrix

$$(H^d_{k_1}(X) + \cdots + H^d_{k_r}(X)) e^{L\log(X)} e^{Q(X)} = H^d(X) e^{L\log(X)} e^{Q(X)}, \tag{2}$$

which is analytic on the sector $\left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left] d - \frac{\pi}{2k_r}, d + \frac{\pi}{2k_r} \right[\right\}$, is a solution of $\partial_X Y(X) = A(X)Y(X)$. As a matter of fact, $H^d_{k_i}(X)$ is k_i-Gevrey asymptotic to $\hat{H}_{k_i}(X)$ on the larger sector $\left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left] d_l - \frac{\pi}{2k_i}, d_{l+1} + \frac{\pi}{2k_i} \right[\right\}$, where d_l, d_{l+1} are two singulars directions and such that $]d_l, d_{l+1}[$ contains no singular directions. Therefore, we can construct an analytic solution on the sector:

$$\left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left] d_l - \frac{\pi}{2k_r}, d_{l+1} + \frac{\pi}{2k_r} \right[\right\}.$$

Let $d \in \mathbb{R}$, and let:

$$d - \frac{\pi}{2k_r} < d^- < d < d^+ < d + \frac{\pi}{2k_r},$$

such that there are no singular directions in $[d^-, d[\bigcup]d, d^+]$. We get two matrices $H^{d^+}(X)e^{L\log(X)}e^{Q(X)}$

and $H^{d^-}(X)e^{L\log(X))}e^{Q(X)}$ which are germs of analytic solutions on the sectors

$$\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in\left]d^- -\frac{\pi}{2k_r},d+\frac{\pi}{2k_r}\right[\right\}\text{ and }\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in\left]d-\frac{\pi}{2k_r},d^+ +\frac{\pi}{2k_r}\right[\right\}.$$

The two matrices are in particular germs of solutions of $\partial_X Y(X)=A(X)Y(X)$ on the sector

$$\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in\left]d-\frac{\pi}{2k_r},d+\frac{\pi}{2k_r}\right[\right\}.$$

A computation shows that there exists a matrix $St_d\in\mathrm{GL}_m(\mathbb{C})$, which we call the Stokes matrix in the direction d, such that:

$$H^{d^+}(X)e^{L\log(X)}e^{Q(X)}=H^{d^-}(X)e^{L\log(X)}e^{Q(X)}St_d.$$

PROPOSITION 1.7. *The following statements are equivalents.*

(i) *The entries of* $\hat{H}(X)$ *converges.*

(ii) $St_d=\mathrm{Id}$ *for all* $d\in\mathbb{R}$.

(iii) $St_d=\mathrm{Id}$ *for all singular directions.*

We can compute the asymptotic solutions using the Laplace and the Borel transformations. See Chapters 2 and 3 of [B] for more details.

DEFINITION 1.8. (1) Let $k\in\mathbb{Q}$. The formal Borel transform $\hat{\mathcal{B}}_k$ is the map that transforms the formal power series $\sum a_nX^n$ into the formal power series:

$$\hat{\mathcal{B}}_k\left(\sum a_nX^n\right)=\sum\frac{a_n}{\Gamma(1+\frac{n}{k})}X^n.$$

(2) Let $d\in\mathbb{R}$, $k\in\mathbb{Q}$, $\varepsilon>0$ and let f analytic on the sector $\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in]d-\varepsilon,d+\varepsilon[\right\}$. We assume that there exists $A,B>0$ such that for $\arg(X)=d$,

$$|f(X)|\leqslant Ae^{B|X|^k}.$$

Then, the following integral is the germ of an analytic function on $\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in\left]d-\frac{\pi}{2k},d+\frac{\pi}{2k}\right[\right\}$ (see [B], page 13 for a proof), and is called the Laplace transform of order k in the direction d of f:

$$\mathcal{L}_{k,d}(f)(X)=\int_0^{\infty e^{id}}f(u)e^{-(\frac{u}{X})^k}d\left(\left(\frac{u}{X}\right)^k\right).$$

For a proof of the following proposition, see Section 7.2 of [B].

PROPOSITION 1.9. *Let* $k_1<\dots<k_r$ *be the levels of* $\partial_X Y(X)=A(X)Y(X)$ *and set* $k_{r+1}=+\infty$. *Suppose that* $d\in\mathbb{R}$ *is not a singular direction, and let* $\hat{h}(X)$ *be an entry of* $\hat{H}(X)$. *Let* $(\kappa_1,\dots,\kappa_r)$ *defined as:*

$$\kappa_i^{-1}=k_i^{-1}-k_{i+1}^{-1}.$$

The series $\hat{\mathcal{B}}_{\kappa_r}\circ\dots\circ\hat{\mathcal{B}}_{\kappa_1}(\hat{h})$ *converges and there exist* $\varepsilon_1,A_1,B_1>0$ *such that it has an analytic continuation* h_1 *on the sector* $\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in]d-\varepsilon_1,d+\varepsilon_1[\right\}$, *and in this sector,*

$$|h_1(X)|\leqslant A_1e^{B_1|X|^{\kappa_1}}.$$

Moreover, for $j=2$ *(resp.* $j=3,\dots,j=r$*), there exist* $\varepsilon_j,A_j,B_j>0$ *such that the function* $h_{j+1}=\mathcal{L}_{\kappa_j,d}(h_j)$ *is analytic on the sector* $\left\{X\in\widetilde{\mathbb{C}}\middle|\arg(X)\in]d-\varepsilon_j,d+\varepsilon_j[\right\}$ *and on this sector*

$$|h_j(X)|\leqslant A_je^{B_j|X|^{\kappa_j}}.$$

Therefore, we may apply $\mathcal{L}_{\kappa_r,d}\circ\cdots\circ\mathcal{L}_{\kappa_1,d}\circ\hat{\mathcal{B}}_{\kappa_r}\circ\cdots\circ\hat{\mathcal{B}}_{\kappa_1}$ to every entries of $\hat{H}(X)$. We have the following equality:

$$H^d(X)=\mathcal{L}_{\kappa_r,d}\circ\cdots\circ\mathcal{L}_{\kappa_1,d}\circ\hat{\mathcal{B}}_{\kappa_r}\circ\cdots\circ\hat{\mathcal{B}}_{\kappa_1}(\hat{H}).$$

1.4 Stokes phenomenon in the parameterized case.

Let $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in \mathrm{M}_m(\mathcal{O}_U)$ (see page 3), where $\varnothing\neq U$ is a polydisc of $\mathbb{C}^n$, and consider $F(X,t)=\hat{H}(X,t)X^{L(t)}e(Q(X,t))$ the fundamental solution of Proposition 1.1. Since for all $k\in\mathbb{N}$, $F(X,t)=\hat{H}(X,t)\mathrm{Diag}(X^k)X^{L(t)-k\mathrm{Id}}e(Q(X,t))$, we may assume that $\hat{H}(X,t)$ has no pole at $X=0$. We want to extend the definition of the singular directions to the parameterized case. Consider $q(X,t)=q_k(t)X^{-k/\nu}+\cdots+q_1(t)X^{-1/\nu}\in\mathbf{E}_U$. A continuous function $d:U\rightarrow\mathbb{R}$ is called singular for $q(X,t)$ if

$$\forall t\in U,\quad q_k(t)e^{-id(t)k/\nu}\in\mathbb{R}^{\geqslant 0}.$$

In general, if $d(t)$ is a singular direction for $q(X,t)$, the positive number $q_k(t)e^{-id(t)k/\nu}$ depends on t. The singular directions of $\partial_X Y(X,t)=A(X,t)Y(X,t)$ (we will write singular directions when no confusion is likely to arise) are the directions that are singular for one of the $q_i(X,t)-q_j(X,t)$, with $i\neq j$. Unfortunately, two different singular directions may be equal on a subset of U. For example, for $n=1$, $U=\mathbb{C}^*$, and $A(X,t)=\mathrm{Diag}(X^{-2},tX^{-2},-tX^{-2})$ we find three exponentials: $e^{-1/X}$, $e^{t/X}$ and $e^{-t/X}$. For $t\in\mathbb{R}^{>0}$, the singular directions of $(2t)X^{-1}$ are the same as singular directions of $(t+1)X^{-1}$. Let $(d_i(t))_{i\in\mathbb{N}}$ be the singular directions, and

$$\mathcal{D}=\left\{t\in U\,|\exists j,j'\in\mathbb{N},\exists t_0\in U,\text{ such that } d_j(t)=d_{j'}(t)\text{ and } d_j(t_0)\neq d_{j'}(t_0)\right\}.$$

Lemma 1.10. *$\mathcal{D}$ is a closed subset of U with empty interior.*

Proof. Assume that there exists a polydisc $\varnothing\neq D\subset\mathcal{D}$, and two singular directions $d_j(t),d_{j'}(t)$, such that $d_j(t)=d_{j'}(t)$ on D. Then, there exists a polydisc $\varnothing\neq D'\subset D$, $q(t),q'(t)\in\mathcal{M}_{D'}$ that do not vanish on D', such that $q(t)/q'(t)$ has constant argument on D'. An analytic function with constant argument on a polydisc is constant. Hence, we deduce that $d_j(t)=d_{j'}(t)$ on a polydisc, which implies that $d_j(t)=d_{j'}(t)$ on U. Since the set of singular directions is finite modulo $2\pi\nu$ with $\nu\in\mathbb{N}^*$, $\mathcal{D}$ has empty interior. □

If we take a smaller polydisc $\varnothing\neq U$, we may assume that $\mathcal{D}=\varnothing$.

From now, we will make the assumption that $\mathcal{D}=\varnothing$.

We still consider $\partial_X Y(X,t)=A(X,t)Y(X,t)$ a parameterized linear differential system with $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$ and $\hat{H}(X,t)X^{L(t)}e(Q(X,t))\in\mathrm{GL}_m\left(\widehat{\mathbf{K}_U}\right)$ the fundamental solution in the same form as in Proposition 1.1. Let $d(t)$ be a singular direction, and let $k_1<\cdots<k_r$ be the levels of $\partial_X Y(X,t)=A(X,t)Y(X,t)$. For t belonging to U, we define the parameterized Stokes matrix $St_{d(t)}$ (we will write Stokes matrix when no confusion is likely to arise) as $t\mapsto St_{d(t)}$, where $St_{d(t)}$ is the Stokes matrix of the specialized system.

Proposition 1.11. *Let $d(t)$ continuous in t, such that for all t_0 in U, $d(t_0)$ is not a singular direction of the unparameterized linear differential equation $\partial_X Y(X,t_0)=A(X,t_0)Y(X,t_0)$. We define $t\mapsto H^{d(t)}(X,t)e^{L(t)\log(X)}e^{Q(X,t)}$, as the solution (2), of the specialized system. Let $d_1(t),d_2(t)$ be two singulars directions such that for all $t\in U$, $d_1(t)<d(t)<d_2(t)$ and $]d_1(t),d_2(t)[$ contains*

no singular directions. Then, there exists a map $U \to \mathbb{R}^{>0}$, $t \mapsto \varepsilon(t)$, *which is not necessary continuous, such that* $H^{d(t)}(X,t)e^{L(t)\log(X)}e^{Q(X,t)}$ *is meromorphic in* (X,t) *for*

$$(X,t) \in \left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]d_1(t) - \frac{\pi}{2k_r}, d_2(t) + \frac{\pi}{2k_r}\right[,\ \text{and } 0 < |X| < \varepsilon(t) \right\} \times U.$$

Proof. We have seen in §1.3, that for t fixed, the asymptotic solution is a germ of meromorphic function on the sector

$$\left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]d_1(t) - \frac{\pi}{2k_r}, d_2(t) + \frac{\pi}{2k_r}\right[\right\}.$$

Since the singular directions are continuous in t, we may assume that $d(t)$ is locally constant. Since for $X \neq 0$, $t \mapsto e^{L(t)\log(X)}e^{Q(X,t)} \in \mathcal{M}_U$, this is now a consequence of Proposition 1.9 and Lemma 1.12 below. □

LEMMA 1.12. *We keep the same notations as in Definition 1.8 and Proposition 1.9. Let* $\hat{h}(X,t)$ *be one of the entries of* $\hat{H}(X,t)$. *Let* $\varnothing \neq V \subset U$ *be a polydisc, and let* $d \in \mathbb{R}$ *such that for all* $t \in V$, d *is not an unparameterized singular direction of* $\partial_X Y(X,t) = A(X,t)Y(X,t)$. *Then, there exists a map* $U \to \mathbb{R}^{>0}$, $t \mapsto \varepsilon(t)$, *which is not necessary continuous, such that*

$$\mathcal{L}_{\kappa_r,d} \circ \cdots \circ \mathcal{L}_{\kappa_1,d} \circ \hat{\mathcal{B}}_{\kappa_r} \circ \cdots \circ \hat{\mathcal{B}}_{\kappa_1}(\hat{h})$$

is meromorphic in (X,t) *on*

$$(X,t) \in \left\{ X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]d - \frac{\pi}{2k_r}, d + \frac{\pi}{2k_r}\right[,\ \text{and } 0 < |X| < \varepsilon(t) \right\} \times V.$$

Moreover, for all $j \leqslant n$:

$$\mathcal{L}_{\kappa_r,d} \circ \cdots \circ \mathcal{L}_{\kappa_1,d} \circ \hat{\mathcal{B}}_{\kappa_r} \circ \cdots \circ \hat{\mathcal{B}}_{\kappa_1}(\partial_{t_j}\hat{h}) = \partial_{t_j}\left(\mathcal{L}_{\kappa_r,d} \circ \cdots \circ \mathcal{L}_{\kappa_1,d} \circ \hat{\mathcal{B}}_{\kappa_r} \circ \cdots \circ \hat{\mathcal{B}}_{\kappa_1}(\hat{h})\right).$$

Proof. We will proceed in two steps.

(1) We recall that $\hat{h}(X,t) \in \hat{K}_U[X^{1/\nu}]$ and (see Remark 1.6) all the X-coefficients are analytics on U. Because of Proposition 1.9, for t fixed, $\hat{\mathcal{B}}_{\kappa_r} \circ \cdots \circ \hat{\mathcal{B}}_{\kappa_1}(\hat{h})$, is a germ of meromorphic function. Therefore, it belongs to $\mathcal{O}_U[X^{1/\nu}]$. Let h_1 be the analytic continuation defined in Proposition 1.9. In particular, for all $X \in \widetilde{\mathbb{C}}$ with $\arg(X) = d$, $t \mapsto h_1(X,t) \in \mathcal{M}_V$. The fact that we have meromorphic function allow us to differentiate termwise and for all $j \leqslant n$, $\partial_{t_j} h_1$ is equals to the analytic continuation of:

$$\hat{\mathcal{B}}_{\kappa_r} \circ \cdots \circ \hat{\mathcal{B}}_{\kappa_1}(\partial_{t_j}\hat{h}).$$

(2) Let $h_2, \dots, h_r$ be the functions defined in Proposition 1.9. Let $t_0 \in V$, let W_{t_0} be a compact neighborhood of t_0 in V, let $i \leqslant r$, and assume that for $X \in \widetilde{\mathbb{C}}$ with $\arg(X) = d$, $t \mapsto h_i(X,t)$ is meromorphic on W_{t_0}. It is sufficient to prove that for all $X \in \widetilde{\mathbb{C}}$ with $\arg(X) \in \left]d - \frac{\pi}{2\kappa_i}, d + \frac{\pi}{2\kappa_i}\right[$ and $|X|$ sufficiently small, $t \mapsto h_{i+1}(X,t)$ is meromorphic on W_{t_0} and for all $j \leqslant n$:

$$\mathcal{L}_{\kappa_i,d}(\partial_{t_j} h_i) = \partial_{t_j}\left(\mathcal{L}_{\kappa_i,d}(h_i)\right) = \partial_{t_j} h_{i+1}.$$

The function $\mathcal{L}_{\kappa_i,d}(h_i)$ is an integral of a meromorphic function depending analytically upon parameters, and we just have to prove that it is possible to find a function f such that, for all $t \in W_{t_0}$, $|h_i(u,t)| < |f(u)|$ and for $\arg(X) \in \left]d - \frac{\pi}{2\kappa_i}, d + \frac{\pi}{2\kappa_i}\right[$, $|X|$ sufficiently small, $\mathcal{L}_{\kappa_i,d}(|f|)(X) < \infty$. From Proposition 1.9, we obtain the existence of $A(t), B(t) > 0$ such that for $\arg(u) = d$ $|h_i(u,t)| \leqslant A(t)e^{B(t)|u|^{\kappa_i}}$. Since $h_i(u,t)$ is meromorphic, we may assume that $A(t)$ and $B(t)$ are continuous on W_{t_0}. The functions $A(t)$ and $B(t)$ admit a maximum A and B on the compact set W_{t_0}. Finally for $\arg(X) \in \left]d-\frac{\pi}{2\kappa_i}, d+\frac{\pi}{2\kappa_i}\right[$ and $|X|$ sufficiently small,

$$|\mathcal{L}_{\kappa_i,d}h_i| = \left|\int_0^{\infty e^{id}} h_i(u,t)e^{-\left(\frac{u}{X}\right)^{\kappa_i}} d\left(\left(\frac{u}{X}\right)^{\kappa_i}\right)\right| \leqslant \int_0^{\infty} Ae^{B|u|^{\kappa_i}} \left|e^{-\left(\frac{u}{X}\right)^{\kappa_i}}\right| d\left(\left(\frac{u}{X}\right)^{\kappa_i}\right) < \infty.$$

□

2. Parameterized differential Galois theory

In this section we are interested to the parameterized differential Galois theory: this is a generalization of the differential Galois theory for parameterized linear differential equations. In §2.1, we review the parameterized differential Galois theory developed in [CS]. In §2.2, we prove that some of the results of §2.1 stay valid without the assumption that the field of constants is differentially closed. This will help us in § 2.3 to prove that the local analytic parameterized differential Galois group descents to a smaller field, whose field of constants is not differentially closed. In §2.4, we explain the main result of the paper: we will show an analogue of the density theorem of Ramis in the parameterized case. In §2.5, we give a similar result for the global parameterized differential Galois group. We end by giving various examples of computation of parameterized differential Galois groups with the parameterized density theorem.

2.1 Basic facts.

We recall some facts from [CS] about Galois theory of parameterized linear differential equations. Classical Galois theory of unparameterized linear differential equation is presented in many books such as [VdPS] and [Ma]. Let K be a differential field of characteristic 0 with $n+1$ commuting derivations: $\partial_0, \dots, \partial_n$. We want to study differential equations of the form $\partial_0 Y = AY$, with $A \in \mathrm{M}_m(K)$. Let C_K be the field of constants with respect to ∂_0. Since all the derivations commute with ∂_0, $(C_K, \partial_1, \dots, \partial_n)$ is a differential field.

Example 2.1. If $K = \hat{K}_U$, then $\partial_0 = \partial_X$, $\{\partial_1, \dots, \partial_n\} = \Delta_t$, and $C_K = \mathcal{M}_U$.

A parameterized Picard-Vessiot extension for the parameterized linear differential equation $\partial_0 Y = AY$ on K is a $(\partial_0, \dots, \partial_n)$-differential field extension $\widetilde{K}|K$ with the following properties:

- There exists a fundamental solution for $\partial_0 Y = AY$ in $\widetilde{K}$ (i.e: an invertible matrix $U = (u_{i,j})$, with entries in $\widetilde{K}$, such that $\partial_0 U = AU$).
- $\widetilde{K} = K\langle u_{i,j}\rangle_{\partial_0,\dots,\partial_n}$ (i.e: $\widetilde{K}$ is the $(\partial_0, \dots, \partial_n)$-differential field generated by K and the $u_{i,j}$).
- The field of constants of $\widetilde{K}$ with respect to ∂_0 is C_K.

Let L be a $(\partial_1, \dots, \partial_n)$-field of characteristic 0 with commuting derivations. The $(\partial_1, \dots, \partial_n)$-differential ring $L\{y_1, \dots, y_k\}_{\partial_1 \dots \partial_n}$ of differential polynomials in k variables over L is the usual polynomial ring in the infinite set of variables

$$\{\partial_1^{\nu_1} \dots \partial_n^{\nu_n} y_j\}_{j \leqslant k}^{\nu_i \in \mathbb{N}},$$

and with derivations extending those in $\{\partial_1, \dots, \partial_n\}$ on L, defined by:

$$\partial_i\left(\partial_1^{\nu_1} \dots \partial_n^{\nu_n} y_j\right) = \partial_1^{\nu_1} \dots \partial_i^{\nu_i+1} \dots \partial_n^{\nu_n} y_j.$$

DEFINITION 2.2 ([CS], Definition 3.2). We say that $(C_K, \partial_1, \dots, \partial_n)$ is differentially closed if it has the following property: For any $k_1, k_2 \in \mathbb{N}$ and for all $P_1, \dots, P_{k_1} \in C_K\{y_1, \dots, y_{k_2}\}_{\partial_1,\dots,\partial_n}$, the system

$$\begin{cases} P_1(\alpha_1, \dots, \alpha_{k_2}) & = & 0 \\ & \vdots & \\ P_{k_1-1}(\alpha_1, \dots, \alpha_{k_2}) & = & 0 \\ P_{k_1}(\alpha_1, \dots, \alpha_{k_2}) & \neq & 0, \end{cases}$$

has a solution in C_K as soon as it has a solution in a $(\partial_1, \dots, \partial_n)$-differential field containing C_K.

For the simplicity of the notations, we will say that C_K differentially closed rather than $(C_K, \partial_1, \dots, \partial_n)$ is differentially closed. Notice that there exists a differentially closed extension of C_K, see [CS], Section 9.1. By definition, a differentially closed field is algebraically closed.

PROPOSITION 2.3 ([CS], Theorem 9.5). *Assume that C_K is differentially closed. Then, we have existence and uniqueness of the parameterized Picard-Vessiot extension for $\partial_0 Y = AY$ up to $(\partial_0, \dots, \partial_n)$-differential isomorphism.*

Until the end of the subsection, we assume that C_K is differentially closed.

The parameterized differential Galois group $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ is the group of field automorphisms of $\widetilde{K}$ which induce the identity on K and commute with all the derivations. In the unparameterized case, the differential Galois group is an algebraic subgroup of $\mathrm{GL}_m(C_K)$. In the parameterized case, we find a linear differential algebraic subgroup:

DEFINITION 2.4. Let us consider m^2 indeterminates $(X_{i,j})_{i,j\leqslant m}$. We say that a subgroup G of $\mathrm{GL}_m(C_K)$ is a linear differential algebraic group if there exist $P_1, \dots, P_k \in C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}$, such that for $A = (a_{i,j}) \in \mathrm{GL}_m(C_K)$,

$$A \in G \Longleftrightarrow P_1(a_{i,j}) = \cdots = P_k(a_{i,j}) = 0.$$

Let U be a fundamental solution of $\partial_0 Y = AY$. We have an injective group morphism:

$$\begin{array}{cccc} \rho_U : & Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right) & \longrightarrow & \mathrm{GL}_m(C_K) \\ & \varphi & \longmapsto & U^{-1}\varphi(U). \end{array}$$

A fundamental fact is that

$$\mathrm{Im}\ \rho_U = \left\{U^{-1}\varphi(U), \varphi \in Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)\right\}$$

is a linear differential algebraic subgroup of $\mathrm{GL}_m(C_K)$ (see Theorem 9.5 in [CS]). If we take a different fundamental solution in $\widetilde{K}$, we obtain a conjugate linear differential algebraic subgroup of $\mathrm{GL}_m(C_K)$. We will identify $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ with a linear differential algebraic subgroup of $\mathrm{GL}_m(C_K)$ for a chosen fundamental solution. We put a topology on $\mathrm{GL}_m(C_K)$, called Kolchin topology, for which the closed sets are defined as the zero loci of finite sets of differential polynomials with coefficients in C_K.

Example 2.5. (Example 3.1 in [CS]) Let $n = 1$, let $\partial_0 = \partial_X$, $\partial_1 = \partial_t$ and let us consider $K = C_K(X)$ the (∂_X, ∂_t)-differential field of rational functions in the indeterminate X, with coefficients in C_K, where X is a ∂_t-constant with $\partial_X X = 1$, C_K is the field of constants with respect to ∂_X, and ∂_X commutes ∂_t. Let us consider the parameterized differential equation

$$\partial_X Y(X,t) = \frac{t}{X} Y(X,t).$$

The fundamental solution is (X^t) and the Parameterized Picard-Vessiot extension is $K(X^t,\log)$ (see §1.1 for the notations). Here, we have added log because we want the extension to be closed under the derivations ∂_X and ∂_t. Using the fact the Galois group commutes with ∂_X and ∂_t, we find that the Galois group is given by:

$$\left\{f\in C_K | f\neq 0 \text{ and } f\partial_t^2 f-(\partial_t f)^2=0\right\}.$$

We can see that if we take $C_K=\mathbb{C}(t)$ or $C_K=\mathcal{M}_{\mathbb{C}}$ (see page 3), which are not differentially closed, then we can find two different groups of differential automorphisms:

$$\left\{f\in \mathbb{C}(t) | f\neq 0 \text{ and } f\partial_t^2 f-(\partial_t f)^2=0\right\}=\mathbb{C}^*$$

and

$$\left\{f\in \mathcal{M}_{\mathbb{C}} | f\neq 0 \text{ and } f\partial_t^2 f-(\partial_t f)^2=0\right\}=\left\{ce^{bt}, b\in\mathbb{C}, c\in\mathbb{C}^*\right\},$$

which shows the importance of considering a Galois group defined over a differentially closed field. See Example 2.23 for the computation of this example using the parameterized density theorem.

We have a Galois correspondence theorem for parameterized differential Galois theory, see Theorem 9.5 in [CS]. For G subgroup of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$, let:

$$\widetilde{K}^G=\left\{a\in\widetilde{K}|\sigma(a)=a,\ \forall\sigma\in G\right\}.$$

The Kolchin closed subgroups of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ are in bijection with the $(\partial_0,\dots,\partial_n)$-differential subfields of $\widetilde{K}$ containing K, via the map:

$$G\mapsto \widetilde{K}^G.$$

The inverse map is given by:

$$M\mapsto Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|M\right),$$

where $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|M\right)$ denotes the set of elements of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ inducing identity on M. In particular, we have the following corollary:

Corollary 2.6. *Let G be a subgroup of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$. Then, $\widetilde{K}^G=K$ if and only if G is dense for Kolchin topology in $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$.*

Let $L|M|K$ be $(\partial_1,\dots,\partial_n)$-differential field extensions. Notice that we do not exclude $L=M=K$. All the definitions we are going to give before the next Proposition are presents in [HS], § 6.2.3.

The $(\partial_1,\dots,\partial_n)$-differential transcendence degree of L over M is the maximum number of elements $a_1,\dots,a_k$ of L such that:

$$P(a_1,\dots,a_k)\neq 0,$$

for all non-zero $(\partial_1,\dots,\partial_n)$-differential polynomials P with coefficients in M.

Let us consider m^2 indeterminates $(X_{i,j})_{i,j\leqslant m}$. Let (p) be a prime $(\partial_1,\dots,\partial_n)$-differential ideal of $C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}$ (i.e: a prime ideal stable under the derivations $\partial_1,\dots,\partial_n$). The $(\partial_1,\dots,\partial_n)$-dimension of (p) over C_K is the $(\partial_1,\dots,\partial_n)$-differential transcendence degree of the quotient ring $C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}/(p)$ over C_K.

Let (r) be a radical $(\partial_1,\dots,\partial_n)$-differential ideal of $C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}$ (i.e: a radical ideal stable under the derivations $\partial_1,\dots,\partial_n$). Let $(p_1),\dots,(p_\nu)$ with $\nu\in\mathbb{N}^*$ be the prime $(\partial_1,\dots,\partial_n)$-differential ideals such that $(r)=\bigcap\limits_{k\leqslant\nu}(p_k)$. The $(\partial_1,\dots,\partial_n)$-dimension of (r) over C_K is the maximum in k of the $(\partial_1,\dots,\partial_n)$-dimension of (p_k) over C_K.

Assume that $M\subset\widetilde{K}$. Let (q) be the radical $(\partial_1,\dots,\partial_n)$-differential ideal of $C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}$ that defines $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|M\right)$ (see the proof of Proposition 9.10 in [CS]). We define the $(\partial_1,\dots,\partial_n)$-differential dimension of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|M\right)$ over C_K as the $(\partial_1,\dots,\partial_n)$-dimension of (q) over C_K.

Proposition 2.7 ([HS], Proposition 6.26). *The $(\partial_1,\dots,\partial_n)$-differential transcendence degree of $\widetilde{K}$ over M is equal to the $(\partial_1,\dots,\partial_n)$-differential dimension of $Gal_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|M\right)$ over C_K.*

Example 2.5 (bis). Let us keep the same notations as in Example 2.5. The parameterized Picard-Vessiot extension is $K(X^t,\log)$ and the Galois group is: $\left\{f\in C_K|f\neq 0 \text{ and } f\partial_t^2 f-(\partial_t f)^2=0\right\}$. The ∂_t-differential dimension of the Galois group is 0 and therefore, X^t satisfies a ∂_t-differential polynomial equation in coefficients in C_K.

2.2 Parameterized differential Galois theory for non-differentially closed field of constants.

Let K be a differential field of characteristic 0 with $n+1$ commuting derivations: $\partial_0,\dots,\partial_n$. Let C_K be the field of constants with respect to ∂_0. Notice that we do not assume that C_K is differentially closed. Let $\partial_0 Y=AY$, with $A\in\mathrm{M}_m(K)$, and assume the existence of $\widetilde{K}|K$, a parameterized Picard-Vessiot extension for $\partial_0 Y=AY$ (see § 2.1). This means in particular that the fields of constants of $\widetilde{K}$ with respect to ∂_0 is C_K. Let $F=(F_{i,j})\in\mathrm{GL}_m\left(\widetilde{K}\right)$ be a fundamental solution such that $\widetilde{K}=K\langle F_{i,j}\rangle_{\partial_0,\dots,\partial_n}$. Let $Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ be the group of $(\partial_0,\dots,\partial_n)$-differential field automorphism of $\widetilde{K}$ letting K invariant.

We extend Definition 2.4 for the field C_K. Let us consider m^2 indeterminates $(X_{i,j})_{i,j\leqslant m}$. We say that a subgroup G of $\mathrm{GL}_m(C_K)$ is a linear differential algebraic group if there exist $P_1,\dots,P_k\in C_K\{X_{i,j}\}_{\partial_1,\dots,\partial_n}$, such that for $A=(a_{i,j})\in\mathrm{GL}_m(C_K)$,

$$A\in G\Longleftrightarrow P_1(a_{i,j})=\cdots=P_k(a_{i,j})=0.$$

The goal of the subsection is the prove:

Proposition 2.8. *(1) Let us consider the injective group morphism:*

$$\begin{array}{cccc}\rho_F: & Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right) & \longrightarrow & \mathrm{GL}_m(C_K)\\ & \varphi & \longmapsto & F^{-1}\varphi(F).\end{array}$$

Then,

$$Im\,\rho_F=\left\{F^{-1}\varphi(F),\varphi\in Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)\right\}$$

is a linear differential algebraic subgroup of $\mathrm{GL}_m(C_K)$. We will identify $Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$ with a linear differential algebraic subgroup of $\mathrm{GL}_m(C_K)$ for a chosen fundamental solution.

(2) Let G be a subgroup of $Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$. If $\widetilde{K}^G=K$, then G is dense for Kolchin topology in $Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$.

Notice that contrary to Corollary 2.6, the converse of (2) is false when C_K is not differentially closed. See [CS], Example 3.1.

Proof. (1) We follow now the proof of Proposition 9.10 in [CS]. We consider the differential polynomial ring:

$$R = K\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n},$$

with the ∂_0-differential structure: $\partial_0(X_{i,j}) = A(X_{i,j})$. Let us consider:

$$S = K\{F_{i,j}, 1/\det(F_{i,j})\}_{\partial_0,\dots,\partial_n},$$

the $(\partial_0,\dots,\partial_n)$-differential ring generated over K by the $F_{i,j}$ and $1/\det(F_{i,j})$. The ring S is an integral domain because it is contained in the $(\partial_0,\dots,\partial_n)$-differential field $\widetilde{K}$. Let q be the prime $(\partial_0,\dots,\partial_n)$-differential ideal such that $R/q \simeq S$. Let $Z_{i,j}$ be the image of $X_{i,j}$ in $S \subset \widetilde{K}$, so $(Z_{i,j})$ is a fundamental solution for $\partial_0 Y = AY$. Consider the following rings:

$$K\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n} \subset \widetilde{K}\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n} = \widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n} \supset C_K\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n},$$

where the indeterminates $Y_{i,j}$ are defined by $(X_{i,j}) = (Z_{i,j})(Y_{i,j})$. We remark that $\partial_0(Y_{i,j}) = 0$. Since we consider fields that are of characteristic 0, the differential ideal:

$$q\widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n} \subset \widetilde{K}\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n} = \widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n},$$

is a radical $(\partial_0,\dots,\partial_n)$-differential ideal (see Corollary A.17 in [VdPS]). The next lemma is an adaptation of Lemma 9.8 in [CS] without the assumption that the field of constants is differentially closed.

Lemma 2.9. *The $(\partial_0,\dots,\partial_n)$-ideal $q\widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$ is generated by:*

$$I = q\widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n} \cap C_K\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}.$$

Proof. Let $(e_i)_{i\in B}$ be a basis of $C_K\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$ over C_K. Let:

$$f = \sum_{i=1}^{n} m_i e_i \in q\widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n},$$

with $m_i \in \widetilde{K}$. By induction on n we will show that $f \in I$. If $n = 0$ or 1 there is nothing to prove. We assume that $n > 1$. We can suppose that $m_1 = 1$ and $m_2 \notin C_K$. Then, because of the fact that the field of constants of $\widetilde{K}$ with respect to ∂_X is C_K:

$$\partial_0(f) = \sum_{i=2}^{n} \partial_0(m_i)e_i \neq 0 \text{ and } f \in q\widetilde{K}\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}.$$

Then, by induction, $\partial_0(f) \in I$. With the same argument:

$$\partial_0(m_2^{-1}f) \in I.$$

Then, $\partial_0(m_2^{-1})f = \partial_0(m_2^{-1}f) - m_2^{-1}\partial_0 f \in I$. Since $\partial_0(m_2^{-1}) \neq 0$, we obtain that $f \in I$. ☐

By Lemma 2.9, $q\widetilde{K}\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}$ is generated by:

$$I = q\widetilde{K}\{X_{i,j}, 1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n} \cap C_K\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}.$$

Clearly I is a $(\partial_1,\dots,\partial_n)$-radical ideal of $C_K\{Y_{i,j}, 1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$. Let $C = (C_{i,j}) \in \mathrm{GL}_m(C_K)$. The following statements are equivalent:

(i) $(C_{i,j}) \in Aut_{\partial_0}^{\partial_1,\dots,\partial_n}\left(\widetilde{K}|K\right)$.

(ii) The map $K\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}\to K\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}$ defined by $(X_{i,j})\mapsto(X_{i,j})(C_{i,j})$ leaves q invariant.

(iii) The map $K\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}\to\widetilde{K}$ defined by $(X_{i,j})\mapsto(Z_{i,j})(C_{i,j})$ sends q to 0.

(iv) The map $\widetilde{K}\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}\to\widetilde{K}$ defined by $(X_{i,j})\mapsto(Z_{i,j})(C_{i,j})$ sends

$$q\widetilde{K}\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}=q\widetilde{K}\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}\text{ to }0.$$

(v) The map $\widetilde{K}\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}\to\widetilde{K}$ defined by $(Y_{i,j})\mapsto(C_{i,j})$ sends

$$q\widetilde{K}\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}\text{ to }0.$$

The theorem is now a consequence of the fact that $q\widetilde{K}\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$ is generated by I, a $(\partial_1,\dots,\partial_n)$-radical ideal of $C_K\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$.

(2) We follow the proof of Proposition 9.10 in [CS], and use the same notations as before. By construction, I of Lemma 2.9 above is the differential ideal that defines the Galois group. Assume that the Kolchin closure of G is not the entire Galois group. Then, there exists $P\in C_K\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}$ such that $P\notin I$ and $P(g)=0$ for all $g\in G$. Lemma 2.9 implies that

$$P\notin J=q\widetilde{K}\{Y_{i,j},1/\det(Y_{i,j})\}_{\partial_1,\dots,\partial_n}.$$

Let

$$T=\{Q\in\widetilde{K}\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}|Q\notin J\text{ and }Q((Z_{i,j})(g_{i,j}))=0,\forall g=(g_{i,j})\in G\}.$$

Since $P\in T$, $T\neq\{0\}$. An element $Q\in T$ can be written as:

$$Q=f_1Q_1+\dots+f_\nu Q_\nu,$$

where $f_i\in\widetilde{K}$ and $Q_i\in K\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}$. Let $Q=f_1Q_1+\dots+f_\nu Q_\nu\in T$ such that:

- $f_1=1$.
- All the f_i are non-zero.
- ν is minimal.

For all $g\in G$, let $Q^g=f_1^gQ_1+\dots+f_\nu^gQ_\nu\in T$. Let $g\in G$. Since $Q-Q^g$ is shorter than Q, and satisfies $(Q-Q^g)\left((Z_{i,j})(g_{i,j})\right)=0$, we have $Q-Q^g\in J$. If $Q-Q^g\neq0$, there exists $l\in\widetilde{K}$ such that $Q-l(Q-Q^g)$ is shorter than Q. Since $Q-l(Q-Q^g)\in T$, this is not possible unless $Q-Q^g=0$. Therefore, $Q=Q^g$, for all $g\in G$, and so $Q\in K\{X_{i,j},1/\det(X_{i,j})\}_{\partial_1,\dots,\partial_n}$. Since $Q(Z_{i,j})=0$, we have $Q\in J$. This yields the result.

□

2.3 A result of descent for the local analytic parameterized differential Galois group.

We keep the notations of Section 1. Let $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$, where $\varnothing\neq U$ is a polydisc of $\mathbb{C}^n$. Note that $\mathcal{O}_U$ is a ring but not a field in general [1] . Since $\mathcal{O}_U\subset\hat{K}_U$, which is a field, $\mathcal{O}_U$ is an integral domain, and we can define K_U as the fraction field of $\mathcal{O}_U$. We have

$$\mathcal{M}_U=\{a\in K_U|\partial_X a=0\}=\{a\in\hat{K}_U|\partial_X a=0\}.$$

---

[1] For example, if $n=1$, $(X-t)^{-1}\notin\mathcal{O}_{\mathbb{C}}$. However, we have $(X-t)^{-1}\in\mathcal{O}_{\mathbb{C}^*}$. More generally let $\alpha(X,t)\in\mathcal{O}_U$. For $t\in U$, let $R(t)$ minimal, such that $|\alpha(X,t)|\neq0$ for $0<|X|<R(t)$. There exists a polydisc $\varnothing\neq U'$ such that there exists $\varepsilon>0$ with $R(t)>\varepsilon$ on U'. In particular, we have $\alpha(X,t)^{-1}\in\mathcal{O}_{U'}$.

Let:

$$F(X,t)=(F_{i,j})=\hat{H}(X,t)X^{L(t)}e(Q(X,t)),$$

be the fundamental solution given in Proposition 1.1. Let us consider $K_U\langle F_{i,j}\rangle_{\partial_X,\Delta_t}=\widetilde{K_U}\subset\widehat{\mathbf{K}_U}$. This field has a field of constants with respect to ∂_X contained in $\overline{\mathcal{M}_U}$, the algebraic closure of $\mathcal{M}_U$, because of Proposition 3.22 in [VdPS]. Since $\overline{\mathcal{M}_U}\cap\widetilde{K_U}=\mathcal{M}_U$, we deduce that $\widetilde{K_U}|K_U$ is a parameterized Picard-Vessiot extension. Therefore, the results of §2.2 may be applied here; and we can define a parameterized differential Galois group $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$, which will be identified with a linear differential algebraic subgroup of $\mathrm{GL}_m(\mathcal{M}_U)$. We want to prove now that it is the "same" as the one of §2.1.

Let C be a (Δ_t)-differentially closed field that contains $\mathcal{M}_U$. Let us define $C[[X]][X^{-1}]$, the (∂_X,Δ_t)-differential field, where X is a (Δ_t)-constant with $\partial_X X=1$, C is the field of constants with respect to ∂_X, and ∂_X commutes with all the derivations. We define the ring $K_U\otimes_{\mathcal{M}_U}C$ with the differential structure given by:

$$\forall a\in K_U,\forall c\in C,\forall\partial\in\{\partial_X,\Delta_t\},\quad \partial(a\otimes_{\mathcal{M}_U}c)=\partial a\otimes_{\mathcal{M}_U}c+a\otimes_{\mathcal{M}_U}\partial c.$$

The (∂_X,Δ_t)-differential ring can be naturally embedded into $C[[X]][X^{-1}]$, which implies that it is an integral domain. Therefore, we may define $\mathcal{K}_{C,U}$, the field of fractions of $K_U\otimes_{\mathcal{M}_U}C$. We see now $\mathcal{K}_{C,U}$ (resp. $K_U\otimes_{\mathcal{M}_U}C$) as a subfield (resp. subring) of $C[[X]][X^{-1}]$.

PROPOSITION 2.10. *Let us keep the same notations. Let* $\partial_X Y(X,t)=A(X,t)Y(X,t)$, *with* $A(X,t)\in\mathrm{M}_m(\mathcal{O}_U)$. *The extension field* $\mathcal{K}_{C,U}\langle F_{i,j}\rangle_{\partial_X,\Delta_t}|\mathcal{K}_{C,U}=\widetilde{\mathcal{K}_{C,U}}|\mathcal{K}_{C,U}$ *is a parameterized Picard-vessiot extension for* $\partial_X Y(X,t)=A(X,t)Y(X,t)$. *Moreover, there exist* $P_1,\dots,P_k\in\mathcal{M}_U\{X_{i,j}\}_{\Delta_t}$, *such that the image of the representation of* $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{K}_{C,U}}|\mathcal{K}_{C,U}\right)$ *(resp.* $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$*) associated to* $F(X,t)$ *is the set of* C*-rational points (resp.* $\mathcal{M}_U$*-rational points) of the the linear differential algebraic subgroup of* $\mathrm{GL}_m(C)$ *(resp.* $\mathrm{GL}_m(\mathcal{M}_U)$*) defined by* $P_1,\dots,P_k$. *More explicitly:*

$$\begin{array}{lcl}\left\{F^{-1}\varphi(F),\varphi\in Gal_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{K}_{C,U}}|\mathcal{K}_{C,U}\right)\right\} & = & \{A=(a_{i,j})\in\mathrm{GL}_m(C)|P_1(a_{i,j})=\dots=P_k(a_{i,j})=0\}\\ \left\{F^{-1}\varphi(F),\varphi\in Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)\right\} & = & \{A=(a_{i,j})\in\mathrm{GL}_m(\mathcal{M}_U)|P_1(a_{i,j})=\dots=P_k(a_{i,j})=0\}.\end{array}$$

Proof. We follow the proof of [MS11], Proposition 3.3. Let (d_k) be a $\mathcal{M}_U$-basis of C. Let us prove that the d_k are linearly independent over $\widetilde{K_U}$. Write $\sum_{k\leqslant\kappa}d_kP_k=0$ with $0\neq P_k\in\widetilde{K_U}$, $\kappa\geqslant 2$ minimal and $P_\kappa=1$. We have $\sum_{k\leqslant\kappa-1}d_k\partial_X P_k=0$. If $\kappa=2$, $\partial_X P_1=0$. If $\kappa>2$, we have that for all k, $\partial_X P_k=0$, because of the minimality of κ. Since $\widetilde{K_U}|K_U$ is a parameterized Picard-vessiot extension, for all k, $P_k\in\mathcal{M}_U$, and the d_k are linearly independent over $\widetilde{K_U}$.

Now, we prove that $\mathcal{K}_{C,U}\langle F_{i,j}\rangle_{\partial_X,\Delta_t}|\mathcal{K}_{C,U}$ is a parameterized Picard-vessiot extension for $\partial_X Y(X,t)=A(X,t)Y(X,t)$. Let $\alpha\in\mathcal{K}_{C,U}\langle F_{i,j}\rangle_{\partial_X,\Delta_t}$ with $\partial_X\alpha=0$. We may assume that $\alpha=\sum d_kP_k$, where $P_k\in\widetilde{K_U}$. We have $\partial_X\alpha=\sum d_k\partial_X P_k=0$. Since the d_k are linearly independent over $\widetilde{K_U}$, we find $\partial_X P_k=0$. Hence, $P_k\in\mathcal{M}_U$, because $\widetilde{K_U}|K_U$ is a parameterized Picard-Vessiot extension. Therefore, $\alpha\in C$ and $\mathcal{K}_{C,U}\langle F_{i,j}\rangle_{\partial_X,\Delta_t}|\mathcal{K}_{C,U}$ is a parameterized Picard-vessiot extension for $\partial_X Y(X,t)=A(X,t)Y(X,t)$.

Let $Y_{i,j}$ be a set of m^2 indeterminates and let I_0, I_1, (∂_X,Δ_t)-differential ideals such that:

$$\begin{array}{lclcl}R_0 & = & K_U\{F_{i,j}\}_{\partial_X,\Delta_t} & = & K_U\{Y_{i,j}\}_{\partial_X,\Delta_t}/I_0\\ R_1 & = & \mathcal{K}_{C,U}\{F_{i,j}\}_{\partial_X,\Delta_t} & = & \mathcal{K}_{C,U}\{Y_{i,j}\}_{\partial_X,\Delta_t}/I_1.\end{array}$$

The group $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$ (resp. $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{K}_{C,U}}|\mathcal{K}_{C,U}\right)$) is the set of $B\in\mathrm{GL}_m(\mathcal{M}_U)$ (resp.

$B\in \mathrm{GL}_m(C)$) such that $(F_{i,j})B$ is again a zero of I_0 (resp. I_1). We just have to prove that $I_1=CI_0$. The inclusion $CI_0\subset I_1$ is clear. Let us prove the other inclusion. Let $P\in I_1$. Without loose of generality, we may assume that $P\in \left(K_U\otimes_{\mathcal{M}_U} C\right)[Y_{i,j}]$. Let us write $P=\sum d_k P_k$, where $P_k\in K_U[Y_{i,j}]$. One finds:

$$P(F_{i,j})=\sum d_k P_k(F_{i,j})=0.$$

Since the d_k are linearly independent over $\widetilde{K_U}$, one finds, $P_k(F_{i,j})=0$, and therefore $I_1=CI_0$.

□

2.4 An analogue of the density theorem in the parameterized case.

Let $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in \mathrm{M}_m(\mathcal{O}_U)$, where $\varnothing\neq U$ is a polydisc of $\mathbb{C}^n$. We want to find topological generators for $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$ for Kolchin topology.

We define now the parameterized monodromy. The notion of monodromy in the unparameterized case is well explained in [VdPS]. For more details about parameterized monodromy, see [CS, MS, Si90].

Definition 2.11. The notations are introduced in § 1.1. We define $\hat{m}$, the formal parameterized monodromy, as follows:

- $\forall \hat{H}(X,t)\in \hat{K}_U, \hat{m}(\hat{H}(X,t))=\hat{H}(X,t)$.
- $\forall a(t)\in \mathcal{M}_U, \hat{m}(X^{a(t)})=e^{2i\pi a(t)}X^{a(t)}$.
- $\hat{m}(\log)=2i\pi+\log$.
- $\forall q(X,t)=\sum a_n X^{-n}\in \mathbf{E}_U=\bigcup\limits_{\nu\in\mathbb{Q}^{>0}} X^{\frac{-1}{\nu}}\mathcal{M}_U[X^{\frac{-1}{\nu}}]$, $\hat{m}(e(q(X,t)))=e(\sum a_n e^{-2i\pi n}X^{-n})$.

From the construction of $\hat{K}_U[\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}]$, $\hat{m}$ induces a well defined (∂_X,Δ_t)-differential ring automorphism of $\hat{K}_U[\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}]$, and then it can be extended as a (∂_X,Δ_t)-differential field automorphism of $\widehat{\mathbf{K}_U}$ letting K_U invariant. Since $\widetilde{K_U}\subset\widehat{\mathbf{K}_U}$, and since $\widetilde{K_U}$ is stable by $\hat{m}$, $\hat{m}$ induces an element of $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$.

Remark 2.12. In the regular singular case with one singularity at 0, the definition of formal parameterized monodromy corresponds to the definition given in [MS].

We now introduce the parameterized exponential torus, which is the subgroup of $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$ consisting of elements that act on the $e(q(X,t))$, with $q(X,t)\in\mathbf{E}_U$.

Definition 2.13. Let α be a character of $\mathbf{E}_U$. We define τ_α as the map

- τ_α is the identity on $\hat{K}_{F,U}$.
- $\forall q(X,t)\in\mathbf{E}_U, \tau_\alpha(e(q(X,t)))=\alpha(q(X,t))e(q(X,t))$.

From the construction of $\hat{K}_U[\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}]$, τ_α induces a well defined (∂_X,Δ_t)-differential ring automorphism of $\hat{K}_U[\log,\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}]$, and then it can be extended to a (∂_X,Δ_t)-differential field automorphism of $\widehat{\mathbf{K}_U}$ letting K_U invariant. Since $\widetilde{K_U}\subset\widehat{\mathbf{K}_U}$, and since $\widetilde{K_U}$ is stable by τ_α, τ_α induces an element of $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$.

The parameterized exponential torus (or simply, the exponential torus) is the subgroup of $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{K_U}|K_U\right)$ consisting of the τ_a, where α is a character of $\mathbf{E}_U$. Notice that the matrices of the exponential torus belongs to $\mathrm{GL}_m(\mathbb{C})$.

Lemma 2.14. *Let $d(t)$ be a singular direction of $\partial_X Y(X,t) = A(X,t)Y(X,t)$ (see* §1.4*). The Stokes matrix $St_{d(t)}$ induces an element of $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{K_U}|K_U\right)$.*

Proof. Let us recall the construction of the Stokes matrices. Let $d(t)$ be a singular direction and let k_r be the biggest level of $\partial_X Y(X,t) = A(X,t)Y(X,t)$. The assumption we have made on $\mathcal{D}$ (see §1.4) tells us that there exists $t \mapsto d^{\pm}(t)$ continuous in t, such that

$$d(t) - \frac{\pi}{2k_r} < d^-(t) < d(t) < d^+(t) < d(t) + \frac{\pi}{2k_r},$$

with no singular directions in $[d^-(t), d(t)[\bigcup]d(t), d^+(t)]$. From the construction of $St_{d(t)}$, and §1.3, we know that

$$H^{d^+(t)}(X,t)e^{L(t)\log(X)}e^{Q(X,t)} = H^{d^-(t)}(X)e^{L(t)\log(X)}e^{Q(X,t)}St_{d(t)}.$$

By construction, the Stokes matrix induces identity on K_U. To prove that the Stokes matrices are elements of $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{K_U}|K_U\right)$, we have to prove that the maps $i^{\pm}$, that send $\hat{H}(X,t)X^{L(t)}e(Q(X,t))$ to $H^{d^{\pm}(t)}(X,t)e^{L(t)\log(X)}e^{Q(X,t)}$, induce (∂_X, Δ_t)-field isomorphisms. From the unparameterized case (see Theorem 2, Section 6.4 of [B]), and the relations satisfied by the symbols log, $\left(X^{a(t)}\right)_{a(t)\in\mathcal{M}_U}$ and $(e(q(X,t)))_{q(X,t)\in\mathbf{E}_U}$ (see §1.1), $i^{\pm}$ induce ∂_X-field isomorphisms.

We want now to prove that if $\hat{H}(X,t)$ admits, $H^{d^{\pm}(t)}(X,t)$ as asymptotic sum in the direction $d^{\pm}(t)$, then $\partial_{t_i}\hat{H}(X,t)$ admits $\partial_{t_i}H^{d^{\pm}(t)}(X,t)$ as asymptotic sum in the direction $d^{\pm}(t)$, for all $i \leqslant n$. This is a consequence of Lemma 1.12 and the fact that we may assume that the $d^{\pm}(t)$ are locally constants. Hence $i^{\pm}$ commute with ∂_{t_i} and $i^{\pm}$ induce (∂_X, Δ_t)-field isomorphisms. ☐

Definition 2.15. Let $d(t)$ be a singular direction of $\partial_X Y(X,t) = A(X,t)Y(X,t)$. The element of $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{K_U}|K_U\right)$ induced by the Stokes matrix in the direction $d(t)$ is the Stokes operator in the direction $d(t)$. For simplicity of notation, we write $St_{d(t)}$ for both Stokes operator and the Stokes matrix in the direction $d(t)$.

Proposition 2.16. *If $\alpha \in \widetilde{K_U}$ is fixed by all the Stokes operators $St_{d(t)}$, the monodromy and the exponential torus, then $\alpha \in K_U$.*

Proof. Let $\overline{\mathcal{M}_U}$ be the algebraic closure of $\mathcal{M}_U$. Proposition 3.25 of [VdPS] implies that if $\alpha \in \widehat{\mathbf{K}_U}$ is fixed by the monodromy and the exponential torus, then $\alpha \in \widehat{\mathbf{K}_U} \cap \overline{\mathcal{M}_U}[[X]][X^{-1}] = \hat{K}_U$. Since $\widetilde{K_U} \subset \widehat{\mathbf{K}_U}$, we have to prove that if $\alpha \in \widetilde{K_U} \cap \hat{K}_U$ is fixed by all the Stokes operators, then $\alpha \in K_U$. Let $\alpha \in \widetilde{K_U} \cap \hat{K}_U$ fixed by all the Stokes operators. Let $F(X,t) = \hat{H}(X,t)X^{L(t)}e(Q(X,t))$ be the fundamental solution defined in Proposition 1.1 and let $(\hat{H}_{i,j})$ be the entries of the matrix $\hat{H}(X,t)$. There exists $P \in K_U\langle X_{i,j}\rangle_{\partial_X,\Delta_t}$, such that $P(\hat{H}_{i,j}) = \alpha$. Let $d(t)$ that satisfies the same properties as in Proposition 1.11. Because of Proposition 1.11, there exists a map $U \to \mathbb{R}^{>0}$, $t \mapsto \varepsilon(t)$, which is not necessary continuous, such that $P(H^{d(t)}_{i,j})$ is meromorphic in (X,t) for

$$(X,t) \in \left\{X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]d_1(t) - \frac{\pi}{2k_r}, d_2(t) + \frac{\pi}{2k_r}\right[\text{ and } 0 < |X| < \varepsilon(t)\right\} \times U,$$

where $d_1(t), d_2(t)$ are two singular directions. Since $P(\hat{H}_{i,j})$ is fixed by all the Stokes operators, $P(H^{d(t)}_{i,j})$ is meromorphic in (X,t) for $0 < |X| < \varepsilon(t)$ and $(X,t) \in \widetilde{\mathbb{C}} \times U$. Moreover,

$P(H_{i,j}^{d(t)})(X,t)=P(H_{i,j}^{d(t)})(e^{2i\pi}X,t)$ on his domain of definition, which means that $P(H_{i,j}^{d(t)})$ is meromorphic in (X,t) for $0<|X|<\varepsilon(t)$ and $(X,t)\in\mathbb{C}\times U$. We recall that K_U consist of function $f(X,t)$ such that for $0<|X|<\varepsilon(t)$, $t\mapsto f(X,t)\in\mathcal{M}_U$. We have, $P(H_{i,j}^{d(t)})\in K_U$. We have seen in Lemma 2.14 that the map that sends $\hat{H}(X,t)X^{L(t)}e(Q(X,t))$ to $H^{d(t)}(X,t)e^{L(t)\log(X)}e^{Q(X,t)}$ induces a (∂_X,Δ_t)-field isomorphism. Since the above map leaves K_U invariant, this implies that $P(\hat{H}_{i,j})\in K_U$. □

We can now prove the main theorem of this paper. We recall some notations. Let $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in \mathrm{M}_m(\mathcal{O}_U)$ (see page 3), let K_U be the fraction field of $\mathcal{O}_U$, and let $\widetilde{K_U}|K_U$ be the parameterized Picard-Vessiot extension defined in the beginning of §2.4. Let $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$ be the field automorphisms of $\widetilde{K_U}$ which commute with all the derivations and leave K_U invariant.

Theorem 2.17 (Parameterized analogue of the density theorem of Ramis). *The group generated by the monodromy, the exponential torus and the Stokes operators is dense for Kolchin topology in* $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$.

Proof. First of all, we have already pointed out that the monodromy, the exponential torus and the Stokes operators are elements of $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$. Using Proposition 2.8, we just have to prove that if $\alpha(X,t)\in\widetilde{K_U}$ is fixed by the monodromy, the exponential torus and the Stokes operators, then it belongs to K_U. This is exactly Proposition 2.16. □

Corollary 2.18. $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|K_U\right)$ *contains a finitely generated Kolchin-dense subgroup.*

Proof. Let $q_1(X,t),\dots,q_\beta(X,t)\in\mathbf{E}_U$, $\mathbb{Q}$-linearly independent, such that

$$\widetilde{K_U}\subset\hat{K}_{F,U}\left(e(q_1(X,t)),\dots,e(q_\beta(X,t))\right).$$

Let τ_i be an element of the exponential torus that fixes the $e(q_j(X,t))$ for $j\neq i$, and that sends $e(q_i(X,t))$ to $ae(q_i(X,t))$, with a not a root of unity.

By the definition of the singular directions (see §1.4), there exists $\nu\in\mathbb{N}^*$ such that the singular directions are finite modulo $2\nu\pi$. Let $d_1(t),\dots,d_k(t)$ be such that, if $d(t)$ is a singular direction, then $d(t)$ is equal to one of the $d_i(t)$ modulo $2\nu\pi$. Let $\alpha(X,t)\in\widetilde{K_U}$ be fixed by the monodromy, $\tau_1,\dots,\tau_\beta$, and $St_{d_1(t)},\dots,St_{d_k(t)}$. Using Proposition 2.8, it is sufficient to prove that $\alpha(X,t)\in K_U$.

We can write $\alpha(X,t)$ as an element of:

$$\hat{K}_{F,U}\left(e(q_1(X,t)),\dots,e(q_{\beta-1}(X,t))\right)(e(q_\beta(X,t))).$$

Since the $q_i(X,t)\in\mathbf{E}_U$ are $\mathbb{Q}$-linearly independent, we know by construction that the $e(Nq_\beta(X,t))$, with $N\in\mathbb{Z}$, are $\mathbb{C}$-linearly independent over $\hat{K}_{F,U}\left(e(q_1(X,t)),\dots,e(q_{\beta-1}(X,t))\right)$. If we add the fact that $\alpha(X,t)$ is fixed by τ_β, we obtain:

$$\alpha(X,t)\in\hat{K}_{F,U}\left(e(q_1(X,t)),\dots,e(q_{\beta-1}(X,t))\right).$$

We apply the same reasoning β times to conclude that $\alpha(X,t)\in\hat{K}_{F,U}\cap\widetilde{K_U}$. By the construction of the Stokes operators, we have that $St_{d(t)}=\mathrm{Id}$ if and only if $St_{2\nu\pi+d(t)}=\mathrm{Id}$, where $\nu\in\mathbb{N}^*$ has been defined in the proof. Proposition 2.16 allows us to conclude that $\alpha(X,t)\in K_U$. □

Remark 2.19. (1) Let $\mathbb{C}(t)\{X\}$ be the subset of $\mathcal{O}_U$ consisting of elements of the form $\sum_{i>N}a_i(t)X^i$, with $a_i(t)\in\mathbb{C}(t)$ and $N\in\mathbb{Z}$. Let us consider $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in\mathrm{M}_m(\mathbb{C}(t)\{X\})$.

Even if we would be able to define a parameterized Picard-Vessiot extension over $\mathbb{C}(t)\{X\}$, we would not have a parameterized analogue of the density theorem of Ramis, because the monodromy is not defined in this case. In general, we have

$$\hat{m}(X^{\alpha(t)})=e^{2i\pi\alpha(t)}X^{\alpha(t)}\notin \mathbb{C}(t)\{X\}(X^{\alpha(t)}).$$

This is why we take a larger field of constants with respect to ∂_X.

(2) Similarly, we can prove that the group generated by the monodromy and the exponential torus is dense for Kolchin topology in $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{K_U}|\hat{K}_U\cap\widetilde{K_U}\right)$.

2.5 The density theorem for the global parameterized differential Galois group.

In this subsection, we consider parameterized linear differential equation of the form:

$$\partial_X Y(X,t)=A(X,t)Y(X,t),$$

with $A(X,t)\in \mathrm{M}_m(\mathcal{M}_U(X))$. We want to prove a density theorem for the global parameterized differential Galois group. The result in the unparameterized case can be found in [M], Proposition 1.3. The parameterized singularities of $\partial_X Y(X,t)=A(X,t)Y(X,t)$ belongs to the algebraic closure of $\mathcal{M}_U$. Because of Remark 1.4, after taking a smaller polydisc $\varnothing\neq U$, we may assume that the set of parameterized singularities belongs to $\mathcal{M}_U$. We will write singularity instead of parameterized singularity when no confusion is likely to arise. Let $S=\{\alpha_1(t),\dots,\alpha_k(t)\}\subset\mathbb{P}_1(\mathcal{M}_U)$ be the set of the singularities of $\partial_X Y(X,t)=A(X,t)Y(X,t)$. For any singularity of $\partial_X Y(X,t)=A(X,t)Y(X,t)$, we may define the set of singular directions of all the singularities by considering

$$\partial_X Y(X-\alpha(t),t)=A(X-\alpha(t),t)Y(X-\alpha(t),t)\text{ if }\infty\not\equiv\alpha(t)\in S$$

and

$$\partial_X Y(X^{-1},t)=A(X^{-1},t)Y(X^{-1},t)\text{ if }\infty\equiv\alpha(t)\in S.$$

Let $(d_{i,j}(t))$ be the singular directions for the singularity $\alpha_i(t)$. As in §1.4, we define:

$$\mathcal{D}_{\alpha_i(t)}=\left\{t\in U\,|\exists j,j'\in\mathbb{N},\exists t_0\in U,\text{ such that }d_{i,j}(t)=d_{i,j'}(t)\text{ and }d_{i,j}(t_0)\neq d_{i,j'}(t_0)\right\}.$$

From Lemma 1.10, all the $\mathcal{D}_{\alpha_i(t)}$ are closed set with empty interior. After taking a smaller polydisc $\varnothing\neq U$, we may assume that:

- There exists $\varepsilon>0$ such that for all $t\in U$ and for all $i,j\leqslant k$:
$$|\alpha_i(t)-\alpha_j(t)|>\varepsilon.$$
- $\mathcal{D}_{\alpha_i(t)}=\varnothing$ for all $i\leqslant k$.

Let $x_0(t)\in\mathcal{M}_U$ and let $\varepsilon>0$ such that:

$$\forall t\in U,\forall i<j\leqslant k,\quad |x_0(t)-\alpha_j(t)|>\varepsilon\text{ and }|\alpha_i(t)-\alpha_j(t)|>\varepsilon.$$

For all $i\leqslant k$ and all $t\in U$, we define $U_{\alpha_i(t)}$, the polydisc in the X-plane with center $\alpha_i(t)$ and with radius ε. Let $d_{\alpha_i}(t)$ be a continuous ray from $\alpha_i(t)$ in $U_{\alpha_i(t)}$, $b_{\alpha_i}(t)$ be the continuous point of $d_{\alpha_i}(t)$ with $|b_{\alpha_i}(t)-\alpha_i(t)|=\varepsilon$ and $\gamma_{\alpha_i}(t)$ be a continuous path in $\mathbb{P}_1(\mathcal{M}_U)$ from $x_0(t)$ to $b_{\alpha_i}(t)$, such that for all $t\in U$ and all $j\leqslant k$, $|\gamma_{\alpha_i}(t)-\alpha_j(t)|>\varepsilon/2$. Analytic continuation of $F(X,t)=(F_{i,j})$, a germ of solution at $x_0(t)$ with the path $\gamma_{\alpha_i}(t)$ and $d_{\alpha_i}(t)$ provides a fundamental solution $F^{d_{\alpha_i}(t)}(X,t)$ on a germ of open sector with vertex $\alpha_i(t)$ bisected by $d_{\alpha_i}(t)$.

Let $\widetilde{\mathcal{M}_U(X)} = \mathcal{M}_U(X)\langle F_{i,j}\rangle_{\partial_X,\Delta_t}$. From the assumptions we have made on $x_0(t)$, we deduce that this field has a field of constants with respect to ∂_X equals to $\mathcal{M}_U$. Therefore, we deduce that $\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)$ is a parameterized Picard-Vessiot extension. The results of §2.2 may be applied here and we can define a parameterized differential Galois group $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$, which will be identified with a linear differential algebraic subgroup of $\mathrm{GL}_m(\mathcal{M}_U)$. As in Proposition 2.10, we want to prove now that it is the "same" as the one of §2.1.

Let C be a (Δ_t)-differentially closed field that contains $\mathcal{M}_U$ and let $C(X)$ denotes the (∂_X, Δ_t)-differential field of rational functions in the indeterminate X, with coefficients in C, where X is a (Δ_t)-constant with $\partial_X X = 1$, C is the field of constants with respect to ∂_X, and ∂_X commutes with all the derivations. With the same reasoning as in Proposition 2.10, we prove:

Proposition 2.20. *Let us keep the same notations. Let* $\partial_X Y(X,t) = A(X,t)Y(X,t)$, *with* $A(X,t) \in \mathrm{M}_m(\mathcal{M}_U(X))$. *The extension field* $C(X)\langle F_{i,j}\rangle_{\partial_X,\Delta_t}|C(X) = \widetilde{C(X)}|C(X)$ *is a parameterized Picard-vessiot extension for* $\partial_X Y(X,t) = A(X,t)Y(X,t)$. *Moreover, there exist* $P_1,\dots,P_k \in \mathcal{M}_U\{X_{i,j}\}_{\Delta_t}$, *such that the image of the representation of* $Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right)$ *(resp.* $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$*) associated to* $F(X,t)$ *is the set of* C*-rational points (resp.* $\mathcal{M}_U$*-rational points) of the the linear differential algebraic subgroup of* $\mathrm{GL}_m(C)$ *(resp.* $\mathrm{GL}_m(\mathcal{M}_U)$*) defined by* $P_1,\dots,P_k$. *More explicitly:*

$$\begin{aligned}
\left\{F^{-1}\varphi(F), \varphi \in Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right)\right\} &= \{A=(a_{i,j}) \in \mathrm{GL}_m(C) | P_1(a_{i,j}) = \cdots = P_k(a_{i,j}) = 0\} \\
\left\{F^{-1}\varphi(F), \varphi \in Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)\right\} &= \{A=(a_{i,j}) \in \mathrm{GL}_m(\mathcal{M}_U) | P_1(a_{i,j}) = \cdots = P_k(a_{i,j}) = 0\}.
\end{aligned}$$

We want to find topological generators for $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ for Kolchin topology.

For $\alpha(t) \in \mathcal{M}_U$, let

$$K_{U,\alpha(t)} = \{f(X-\alpha(t),t) \mid f(X,t) \in K_U\},$$

and let

$$K_{U,\infty} = \{f(X^{-1},t) \mid f(X,t) \in K_U\}.$$

Let $\alpha(t) \in S$ and let $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,\alpha(t)} \cap \widetilde{\mathcal{M}_U(X)}\right)$ be the local Galois group for the fundamental solution $F^{d_\alpha(t)}(X,t)$ described above. If we "conjugate" $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,\alpha(t)} \cap \widetilde{\mathcal{M}_U(X)}\right)$ by the analytic continuation of $F(X,t)$ described above, we get an injective morphism of linear differential algebraic group:

$$Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,\alpha(t)} \cap \widetilde{\mathcal{M}_U(X)}\right) \hookrightarrow Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right).$$

Using the maps $i^{\pm}$ defined in the proof of Lemma 2.14 and the injection above, we can define the monodromy, the exponential torus, and the Stokes operators for any singularities in S, as elements of

$$Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right).$$

Theorem 2.21. *Let* $\partial_X Y(X,t) = A(X,t)Y(X,t)$*, where* $A(X,t) \in \mathrm{M}_m(\mathcal{M}_U(X))$*. For* $\alpha(t) \in S$*, let* $G_{\alpha(t)}$ *be the subgroup of:*

$$Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,\alpha(t)} \cap \widetilde{\mathcal{M}_U(X)}\right),$$

generated by the monodromy, the exponential torus and the Stokes operators. Let G *be the subgroup of* $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ *generated by the* $G_{\alpha(t)}$*, with* $\alpha(t) \in S$*. Then* G *in dense for Kolchin topology in*

$$Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right).$$

Proof. We use Proposition 2.8. We have to prove that the subfield of $\widetilde{\mathcal{M}_U(X)}$ fixed by G is $\mathcal{M}_U(X)$. Let $f(X,t) \in \widetilde{\mathcal{M}_U(X)}$ be fixed by G. Then, by the same reasoning as in Proposition 2.16, it follows that $f(X,t)$ belongs to $K_{U,\alpha(t)}$, for $\alpha(t) \in S$. Therefore, we deduce that $f(X,t)$ is meromorphic in (X,t) on $\mathbb{P}_1(\mathbb{C}) \times U$, and has a finite number of poles in the X-plane for t fixed. Hence, $f(X,t) \in \mathcal{M}_U(X)$. □

In particular, we generalize Theorem 4.2 in [MS11] who says that if the equation has only regular singular poles, then the group generated by the monodromy at each pole is dense for Kolchin topology in $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$.

Corollary 2.22. $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ *contains a finitely generated Kolchin-dense subgroup.*

Proof. In the proof of Theorem 2.21, we see that the global parameterized differential Galois group is generated by all local parameterized differential Galois groups. Since there is a finite number of singularities, this is a consequence of Corollary 2.18. □

2.6 Examples.

In all the examples, we will compute the global parameterized differential Galois group. This means that the base field is $\mathcal{M}_U(X)$.

Example 2.23. Let us consider $\partial_X Y(X,t) = \frac{t}{X}Y(X,t)$. This example was treated by hand in Example 2.5 but we will compute here $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ using the parameterized density theorem. The fundamental solution is (X^t) and the parameterized Picard-Vessiot extension over $\mathcal{M}_U(X)$ is $\mathcal{M}_U(X, X^t, \log)$ (we want the extension to be closed under the derivations ∂_X and ∂_t). The exponential torus and the Stokes matrices are the identity. The monodromy sends X^t to $e^{2i\pi t}X^t$. The element $e^{2i\pi t}$ satisfies the differential equation

$$\partial_t\left(\frac{\partial_t e^{2i\pi t}}{e^{2i\pi t}}\right) = 0.$$

Therefore, the Kolchin closure of the monodromy is contained in:

$$\left\{a \in \mathcal{M}_U \middle| \partial_t\left(\frac{\partial_t a}{a}\right)\right\} = \{ce^{bt}, b \in \mathbb{C}, c \in \mathbb{C}^*\}.$$

Conversely, the map that sends X^t to $ce^{bt}X^t$ is an element of $Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$. Finally, viewed as a linear differential algebraic subgroup of $\mathrm{GL}_1(\mathcal{M}_U)$,

$$\begin{array}{rcl} Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right) & \simeq & \left\{a\in\mathcal{M}_U|\partial_t\left(\frac{\partial_t a}{a}\right)=0\right\} \\ & = & \{a\in\mathcal{M}_U|a\neq 0 \text{ and } a\partial_t^2 a-(\partial_t a)^2=0\} \\ & \subseteq & \mathrm{GL}_1(\mathcal{M}_U). \end{array}$$

Example 2.24 (Parameterized Euler equation). Let us consider:

$$\begin{pmatrix}\partial_X Y(X,t)\\ \partial_X^2 Y(X,t)\end{pmatrix}=\begin{pmatrix}0 & 1\\ \frac{-1}{f(t)X^3} & \frac{1}{f(t)X^2}-\frac{1}{X}\end{pmatrix}\begin{pmatrix}Y(X,t)\\ \partial_X Y(X,t)\end{pmatrix},$$

where $f(t)$ is a meromorphic function different from 0. A fundamental solution is:

$$\begin{pmatrix}1 & \hat{F}(X,t)\\ \frac{1}{f(t)X^2} & \partial_X\hat{F}(X,t)\end{pmatrix}\begin{pmatrix}e\left(\frac{-1}{f(t)X}\right) & 0\\ 0 & 1\end{pmatrix},$$

where $\hat{F}(X,t)=\sum\limits_{n\geqslant 0} n!(f(t)X)^{n+1}$. The only singularity is 0. The monodromy is trivial. Let τ be an element of the exponential torus. Then, the image of the fundamental solution under τ is:

$$\begin{pmatrix}1 & \hat{F}(X,t)\\ \frac{1}{f(t)X^2} & \partial_X\hat{F}(X,t)\end{pmatrix}\begin{pmatrix}\alpha e\left(\frac{-1}{f(t)X}\right) & 0\\ 0 & 1\end{pmatrix},$$

with $\alpha\in\mathbb{C}^*$. Therefore, the matrices of the elements of the exponential torus are of the form $\mathrm{Diag}(\alpha,1)$, with $\alpha\in\mathbb{C}^*$. The only level of the system is 1 and the singular directions are the $\arg(f(t))^{-1}+2k\pi$, with $k\in\mathbb{Z}$. As we have seen in Proposition 1.9, we can compute the Stokes matrix with the Laplace and the Borel transforms. To compute the matrix in the direction $\arg(f(t))^{-1}$, we will compute:

$$\mathcal{L}_{1,\arg(f(t))^{-1}+\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t)))-\mathcal{L}_{1,\arg(f(t))^{-1}-\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t))),$$

where $0<\varepsilon<\frac{\pi}{2}$ is taken such that there are no singular directions on the sector:

$$\left[\arg(f(t))^{-1}-\varepsilon,\arg(f(t))^{-1}\right[\quad\bigcup\quad\left]\arg(f(t))^{-1},\arg(f(t))^{-1}+\varepsilon\right].$$

It follows from the definition of the formal Borel transform that $\hat{\mathcal{B}}(\hat{F}(X,t))=\log(1-f(t)X)$. Finally,

$$\begin{array}{rc} & \mathcal{L}_{1,\arg(f(t))^{-1}+\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t)))-\mathcal{L}_{1,\arg(f(t))^{-1}-\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t))) \\ = & X^{-1}\int_0^{\infty i(\arg(f(t))^{-1}+\varepsilon)}\log(1-f(t)u)e^{-(\frac{u}{X})}d(u) \\ - & X^{-1}\int_0^{\infty i(\arg(f(t))^{-1}-\varepsilon)}\log(1-f(t)u)e^{-(\frac{u}{X})}d(u). \end{array}$$

Integration by parts and residue theorem imply that:

$$\mathcal{L}_{1,\arg(f(t))^{-1}+\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t)))-\mathcal{L}_{1,\arg(f(t))^{-1}-\varepsilon}(\hat{\mathcal{B}}(\hat{F}(X,t)))=2i\pi f(t)e^{-\left(\frac{1}{f(t)X}\right)}.$$

Therefore, the Stokes matrix in this direction is $\begin{pmatrix} 1 & 2i\pi f(t) \\ 0 & 1 \end{pmatrix}$. Finally we obtain:

$$\begin{aligned} Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right) &\simeq \left\{\begin{pmatrix} \alpha & bf \\ 0 & 1 \end{pmatrix}, \text{ where } \alpha \in \mathbb{C}^* \text{ and } b \in \mathbb{C}\right\} \\ &\simeq \left\{\begin{pmatrix} \alpha & \beta \\ 0 & 1 \end{pmatrix}, \text{ where } \partial_t \alpha = 0, \alpha \neq 0 \text{ and } \partial_t\left(\frac{\beta}{f}\right) = 0\right\}. \end{aligned}$$

Example 2.25 (Bessel equation). We are interested in the parameterized linear differential equation:

$$\begin{pmatrix} \partial_X Y(X,t) \\ \partial_X^2 Y(X,t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \frac{(t^2-X^2)}{X^2} & \frac{-1}{X} \end{pmatrix} \begin{pmatrix} Y(X,t) \\ \partial_X Y(X,t) \end{pmatrix}.$$

This equation has two singularities: 0 and ∞. Let $\varnothing \neq U$ be a polydisc such that $U \cap (1/2+\mathbb{Z}) = \varnothing$. First, we will compute:

$$Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,0} \cap \widetilde{\mathcal{M}_U(X)}\right).$$

If $t+1/2 \notin \mathbb{Z}$, the two solutions:

$$J_t(X) = \left(\frac{X}{2}\right)^t \sum_{k=0}^{\infty} \frac{(-1)^k X^{2k}}{k!\Gamma(t+k+1)2^k}$$

$$J_{-t}(X) = \left(\frac{X}{2}\right)^{-t} \sum_{k=0}^{\infty} \frac{(-1)^k X^{2k}}{k!\Gamma(-t+k+1)2^k},$$

are linearly independent (see [Wat] Page 43) and we have a fundamental solution. The equation is regular singular, therefore, the group generated by the monodromy is dense for Kolchin topology in the parameterized differential Galois group $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,0} \cap \widetilde{\mathcal{M}_U(X)}\right)$. By the same reasoning as in Example 2.23:

$$Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,0} \cap \widetilde{\mathcal{M}_U(X)}\right) \simeq \left\{\begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix}, \text{ where } \alpha \neq 0 \text{ and } \alpha\partial_t^2\alpha - (\partial_t\alpha)^2 = 0\right\}.$$

We turn now to the singularity at infinity. We have:

$$\begin{pmatrix} \partial_X Y(X^{-1},t) \\ \partial_X^2 Y(X^{-1},t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \frac{t^2}{X^2} - \frac{1}{X^4} & \frac{-1}{X} \end{pmatrix} \begin{pmatrix} Y(X^{-1},t) \\ \partial_X Y(X^{-1},t) \end{pmatrix}.$$

In order to compute the matrices of the monodromy, the elements of the exponential torus, and the Stokes operators, we will help ourself with another basis of solutions:

$$H_t^1(X^{-1}) = \frac{J_{-t}(X^{-1}) - e^{-it\pi} J_t(X^{-1})}{i\sin(t\pi)}$$

$$H_t^2(X^{-1}) = \frac{J_{-t}(X^{-1}) - e^{it\pi} J_t(X^{-1})}{-i\sin(t\pi)}.$$

In [Wat] page 198, we find that on the sector $]-\pi, 2\pi[$, $H_t^1(X^{-1})$ is asymptotic to:

$$\widetilde{H}_t^1(X^{-1}) = \left(\frac{2X}{\pi}\right)^{1/2} e^{i(X^{-1}-t\pi/2-\pi/4)} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(t+k+1/2) X^k}{(2i)^k k! \Gamma(t-k+1/2)}.$$

The same holds for $H_t^2(X^{-1})$ on the sector $]-2\pi,\pi[$:

$$\widetilde{H}_t^2(X^{-1}) = \left(\frac{2X}{\pi}\right)^{1/2} e^{-i(X^{-1}-t\pi/2-\pi/4)} \sum_{k=0}^{\infty} \frac{\Gamma(t+k+1/2)X^k}{(2i)^k k!\Gamma(t-k+1/2)}.$$

It follows that in the basis $(H_t^1(X^{-1}), H_t^2(X^{-1}))$, the matrix of the monodromy is:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

and the matrices of the elements of the exponential torus are of the form:

$$\left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix}, \text{ where } \alpha \in \mathbb{C}^* \right\}.$$

The only level is 1 and due to the expression of $\widetilde{H}_t^1(X^{-1})$ and $\widetilde{H}_t^2(X^{-1})$, the singular directions are the directions $\frac{\pi}{2}+k\pi$, with $k \in \mathbb{Z}$. By definition, the Stokes matrix in the direction $\frac{\pi}{2}+k\pi$ is the matrix that sends the asymptotic representation defined on the sector $](k-1)\pi,(k+1)\pi[$ to the asymptotic representation defined on the sector $]k\pi,(k+2)\pi[$. In [RM1], 3.4.12, we find that in the basis $(H_t^1(X^{-1}), H_t^2(X^{-1}))$ the Stokes matrix in the direction $\frac{\pi}{2}+2k\pi$ is

$$\begin{pmatrix} 1 & 0 \\ 2e^{2i\pi t}\cos(\pi t) & 1 \end{pmatrix},$$

and the Stokes matrix in the direction $-\frac{\pi}{2}+2k\pi$ is

$$\begin{pmatrix} 1 & -2e^{-2i\pi t}\cos(\pi t) \\ 0 & 1 \end{pmatrix}.$$

An application of Theorem 2.21 and Theorem 2.17 gives that $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|K_{U,\infty}\cap\widetilde{\mathcal{M}_U(X)}\right)$ and $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ are linear differential algebraic subgroups of $\mathrm{SL}_2(\mathcal{M}_U)$, because all matrices we have computed are in $\mathrm{SL}_2(\mathcal{M}_U)$, and the Kolchin closure of a subgroup of $\mathrm{SL}_2(\mathcal{M}_U)$ is a linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_U)$.

Let C be a differentially closed field that contains $\mathcal{M}_U$ and consider $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{C(X)}|C(X)\right)$, the parameterized differential Galois group defined in Proposition 2.20. We are going now to compute the Zariski closure G of $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{C(X)}|C(X)\right)$. Let $C^* = C\setminus\{0\}$. From the classification of linear algebraic subgroup of $\mathrm{SL}_2(C)$ (see [VdPS], Theorem 4.29), there are four possibilities:

(i) G is conjugate to a subgroup of $B = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}, \text{ where } a \in C^*, b \in \mathbb{C} \right\}$.

(ii) G is conjugate to a subgroup of $D_\infty = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \bigcup \begin{pmatrix} 0 & b^{-1} \\ -b & 0 \end{pmatrix}, \text{ where } a,b \in C^* \right\}$.

(iii) G is finite.

(iv) $G = \mathrm{SL}_2(\mathrm{C})$.

From Proposition 2.20, every matrices that belongs to $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ belongs also to $Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right)$. Since G must contain

$$\begin{pmatrix} 1 & 0 \\ 2e^{2i\pi t}\cos(\pi t) & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & -2e^{-2i\pi t}\cos(\pi t) \\ 0 & 1 \end{pmatrix},$$

we find that the only possibility is $G = \mathrm{SL}_2(\mathrm{C})$. In [C72], Proposition 42, Cassidy classifies the Zariski-dense differential algebraic subgroups of $\mathrm{SL}_2(C)$. Finally, we have two possibilities:

- $Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right)$ is conjugate to $\mathrm{SL}_2(C_0)$ over $\mathrm{SL}_2(C)$, where $C_0 = \{a \in C(X)|\partial_X a = \partial_t a = 0\}$.
- $Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right) = \mathrm{SL}_2(C)$.

If $Gal^{\Delta_t}_{\partial_X}\left(\widetilde{C(X)}|C(X)\right)$ is conjugate to $\mathrm{SL}_2(C_0)$ over $\mathrm{SL}_2(C)$, the matrix of the monodromy of the singularity 0 is conjugate to a matrix $M \in \mathrm{SL}_2(C_0)$ over $\mathrm{SL}_2(C)$. Similar matrices have the same eigenvalues, then the eigenvalues of M are $e^{2i\pi t}$ and $e^{-2i\pi t}$, which is not possible if M belongs to $\mathrm{SL}_2(C_0)$. Because of Proposition 2.20, we find that

$$Aut^{\Delta_t}_{\partial_X}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right) = \mathrm{SL}_2(\mathcal{M}_U).$$

3. Applications.

We now give three applications of the parameterized differential Galois theory. In §3.1, we deal with linear differential equations that are completely integrable. It has been proved in [CS] that the equation is completely integrable if and only if the parameterized differential Galois group is conjugate over a differentially closed field to a group of constant matrices. We use Theorem 2.21 to prove that the equation is completely integrable if and only if there exists a fundamental solution such that the matrices of the topological generators for the Galois group appearing in Theorem 2.21 are constant matrices. As a corollary, we prove that the equation is completely integrable if and only if the matrices of the topological generators for the Galois group given in the parameterized density theorem are conjugate over $\mathrm{GL}_m(\mathcal{M}_U)$ to constant matrices. Notice that $\mathcal{M}_U$ is not differentially closed. In §3.2, we study an entry of a Stokes operators at the singularity at infinity of the equation:

$$\partial_X^2 Y(X,t) = (X^3 + t)Y(X,t).$$

In particular, we prove that it is not ∂_t-finite: it satisfies no parameterized linear differential equation. This partially answers a question by Sibuya. In §3.3, we deal with the inverse problem in the parameterized differential Galois theory. Let k be a so-called universal (Δ_t)-field, see §3.2. We give a necessary condition on a linear differential algebraic subgroup of $\mathrm{GL}_m(k)$ for being the global parameterized differential Galois group for some equation having coefficients in $k(X)$. The sufficient condition has been proved in [MS11], Corollary 5.2.

3.1 Completely integrable equations.

In this subsection, we study completely integrable equations. See also [GO] for an approach from the point of view of differential Tannakian categories.

Definition 3.1. Let $A_0 \in \mathrm{M}_m(\mathcal{M}_U(X))$. We say that the linear differential equation $\partial_0 Y = A_0 Y$

is completely integrable if there exist $A_1,\dots,A_n \in \mathrm{M}_m(\mathcal{M}_U(X))$ such that, for all $0 \leqslant i,j \leqslant n$,

$$\partial_{t_i} A_j - \partial_{t_j} A_i = A_i A_j - A_j A_i,$$

with $\partial_{t_0} = \partial_X$.

Remark 3.2. Sibuya shows in [Si90], Theorem A.5.2.3, that if the parameterized linear differential equation $\partial_X Y(X,t) = A(X,t)Y(X,t)$ is regular singular, then it is isomonodromic (see page 2 for the definition) if and only if it is completely integrable. This result is not true in the irregular case. The main reason is the fact that there are more topological generators in the parameterized differential Galois group, see Proposition 3.3.

PROPOSITION 3.3. *Let* $A_0(X,t) \in \mathrm{M}_m(\mathcal{M}_U(X))$ *and let* $\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)$ *be the parameterized Picard-Vessiot extension for* $\partial_X Y(X,t) = A_0(X,t)Y(X,t)$ *defined in* §2.5. *The linear differential equation* $\partial_X Y(X,t) = A_0(X,t)Y(X,t)$ *is completely integrable if and only if there exists a fundamental solution* $F(X,t)$ *in* $\widetilde{\mathcal{M}_U(X)}$*, such that the images of the topological generators of* $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ *(see Theorem 2.21), with respect to the representation associated to* $F(X,t)$*, belongs to* $\mathrm{GL}_m(\mathbb{C})$.

Proof. Let C be a differentially closed field that contains $\mathcal{M}_U$ and let us consider $C(X)$ as in §2.5. Let $\widetilde{C(X)}|C(X)$ be the parameterized Picard-Vessiot extension for $\partial_X Y(X,t) = A_0(X,t)Y(X,t)$, and let $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{C(X)}|C(X)\right)$ be the parameterized differential Galois group defined in §2.1. We recall that if we take a different fundamental solution in $\widetilde{\mathcal{M}_U(X)}$ to compute the Galois group, we obtain a conjugate linear differential algebraic subgroup of $\mathrm{GL}_m(\mathcal{M}_U)$.

Using Theorem 2.21, we find that there exists a fundamental solution such that the matrices of the topological generators for the Galois group appearing in Theorem 2.21 are constant if and only if $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ is conjugate over $\mathrm{GL}_m(\mathcal{M}_U)$ to a subgroup of $\mathrm{GL}_m(\mathbb{C})$. Using Proposition 2.20, we find that $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ is conjugate over $\mathrm{GL}_m(\mathcal{M}_U)$ to a subgroup of $\mathrm{GL}_m(\mathbb{C})$ if and only if $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{C(X)}|C(X)\right)$ is conjugate over $\mathrm{GL}_m(C)$ to a subgroup of $\mathrm{GL}_m(C_0)$, where

$$C_0 = \{a \in C(X) | \partial_X a = \partial_{t_1} a = \cdots = \partial_{t_n} a = 0\}.$$

Proposition 3.9, [CS] says that this occurs if and only if there exist $A_1,\dots,A_n \in \mathrm{M}_m(C(X))$ such that, for all $0 \leqslant i,j \leqslant n$,

$$\partial_{t_i} A_j - \partial_{t_j} A_i = A_i A_j - A_j A_i,$$

with $\partial_{t_0} = \partial_X$. To finish, we follow the proof of Proposition 1.24 in [DVH]. Let $0 < i \leqslant n$ and let us consider

$$\partial_X A_i - \partial_{t_i} A_0 = A_0 A_i - A_i A_0.$$

By clearing the denominators, we obtain that every entries of every X-coefficients of A_i satisfies a finite set of polynomial equations in coefficients in $\mathcal{M}_U$. Since the polynomial equations have a solution in C, it must have a solution in the algebraic closure of $\mathcal{M}_U$. Using Remark 1.4, we find the existence of $\varnothing \neq U' \subset U$, such that all the A_i belongs to $\mathrm{M}_m(\mathcal{M}_{U'}(X))$. This concludes the proof. □

In the proof of Proposition 3.3, we have proved:

Corollary 3.4. *Let* $A(X,t)\in \mathrm{M}_m(\mathcal{M}_U(X))$. *The equation* $\partial_X Y(X,t)=A(X,t)Y(X,t)$, *is completely integrable if and only if the matrices of the topological generators for the Galois group appearing in Theorem 2.21 are conjugate over* $\mathrm{GL}_m(\mathcal{M}_U)$ *to constant matrices.*

Remark 3.5. This corollary improves Proposition 3.9 in [CS]. The conjugation occurs in a field that is not differentially closed. Furthermore, we do not need for the entire parameterized differential Galois group to be conjugate to a group of constant matrices in order to deduce that the equation $\partial_X Y(X,t)=A(X,t)Y(X,t)$ is completely integrable.

3.2 On the hyper transcendence of a Stokes matrix.

In this subsection, we will study the parameterized linear differential equation:

$$\partial_X^2 Y(X,t)=(X^3+t)Y(X,t). \tag{3}$$

Sibuya, in Chapter 2 of [Si], shows that there exists a solution $y_0(X,t)$ which admits an asymptotic representation $\widetilde{y}_0(X,t)$ on the sector (see Theorem 6.1 in [Si]):

$$\left\{X\in\widetilde{\mathbb{C}}\,\middle|\,\arg(X)\in\left]\frac{-3\pi}{5},\frac{3\pi}{5}\right[\right\}.$$

We easily check that

$$y_k(X,t)=y_0\left(e^{\frac{-2ki\pi}{5}}X,e^{\frac{-6ki\pi}{5}}t\right)$$

are solutions of (3) which have the asymptotic representation $\widetilde{y}_k(X,t)=\widetilde{y}_0(e^{\frac{-2ki\pi}{5}}X,e^{\frac{-6ki\pi}{5}}t)$ on the sector $S_{k-1}\cup\bar{S}_k\cup S_{k+1}$, where

$$S_k=\left\{X\in\widetilde{\mathbb{C}}\,\middle|\,\arg(X)\in\left]\frac{(2k-1)\pi}{5},\frac{(2k+1)\pi}{5}\right[\right\},$$

and $\bar{S}_k$ is its closure.



The sectors S_k



The singular directions

The asymptotic representation $\widetilde{y}_k(X,t)$ is bounded uniformly on each compact set in the t-plane as $|X|$ tends to infinity on the sector S_k, and tends to infinity uniformly on each compact set in the t-plane as $|X|$ tends to infinity on the sectors S_{k-1} and S_{k+1}. As we see in [Si], page 83, $y_{k+1}(X,t)$ and $y_{k+2}(X,t)$ are linearly independent and we can write $y_k(X,t)$ as a $\mathcal{M}_{\mathbb{C}}$-linear combination of $y_{k+1}(X,t)$ and $y_{k+2}(X,t)$:

$$\forall k\in\mathbb{N},\forall X,t\in\mathbb{C},\ y_k(X,t)=C_k(t)y_{k+1}(X,t)+\widetilde{C}_k(t)y_{k+2}(X,t),\tag{4}$$

where $\widetilde{C}_k(t),C_k(t)\in\mathcal{M}_{\mathbb{C}}$. By Theorem 21.1 in [Si], we obtain that

$$\widetilde{C}_k(t)=-e^{\frac{2i\pi}{5}}\text{ and }C_k(t)=C_0\left(e^{\frac{-6ki\pi}{5}}t\right).$$

In [Si], the author asks if $C_0(t)$ is differentially transcendental (i.e: satisfies no differential polynomial equations). We will use Galois theory to prove that for all polydisc $U\neq\varnothing$, $C_0(t)$ is not ∂_t-finite over $\mathcal{M}_U$ (i.e: satisfies no linear differential equations in coefficients in $\mathcal{M}_U$).
The singularity of the system is at infinity. Let $W(X,t)=XY(X^{-1},t)$. We obtain the parameterized linear differential equation

$$X^7\partial_X^2W(X,t)=(1+tX^3)W(X,t),\tag{5}$$

which can be written in the form

$$\begin{pmatrix}\partial_XW(X,t)\\ \partial_X^2W(X,t)\end{pmatrix}=\begin{pmatrix}0&1\\ \frac{1+tX^3}{X^7}&0\end{pmatrix}\begin{pmatrix}W(X,t)\\ \partial_XW(X,t)\end{pmatrix}.$$

Let k be a so-called universal (Δ_t)-field of characteristic 0: for any (Δ_t)-field $k_0\subset k$, (Δ_t)-finitely generated over $\mathbb{Q}$, and any (Δ_t)-finitely generated extension k_1 of k_0, there is a (Δ_t)-differential k_0-isomorphism of k_1 into k. See Chapter 3, Section 7 of [Kol76] for more details. In particular, k is (Δ_t)-differentially closed. Let $k(X)$ denotes the (∂_X,Δ_t)-differential field of rational functions in the indeterminate X, with coefficients in k, where X is a (Δ_t)-constant with $\partial_XX=1$, k is the field of constants with respect to ∂_X, and ∂_X commutes with all the derivations.

Let $A(X,t)=\begin{pmatrix}0&1\\ \frac{1+tX^3}{X^7}&0\end{pmatrix}$. The two solutions $Xy_1(X^{-1},t)$, $Xy_2(X^{-1},t)$ admit asymptotic representation and the only singularity is 0. Therefore,

$$\mathcal{M}_U(X)\langle y_1(X^{-1},t),y_2(X^{-1},t)\rangle_{\partial_X,\partial_t}|\mathcal{M}_U(X)=\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)$$

is a parameterized Picard-Vessiot extension for $\partial_XW(X,t)=A(X,t)W(X,t)$. Because of Proposition 2.20, $\widetilde{k(X)}|k(X)=k(X)\langle y_1(X^{-1},t),y_2(X^{-1},t)\rangle_{\partial_X,\partial_t}|k(X)$ is a parameterized Picard-Vessiot extension.

LEMMA 3.6. $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)=\mathrm{SL}_2(k)$.

Notice that the differential equation is of the form $\partial_X^2W(X,t)=r(X,t)W(X,t)$, where $r(X,t)\in k(X)$. In this case, we can compute the Galois group using a parameterized version of Kovacic's algorithm, see [Ar, D]. In order to have a self contain proof, we will perform the computations explicitly.

Proof. If we apply Kovacic's algorithm (see [Kov]), we find that the unparameterized differential Galois group $Gal_{\partial_X}\left(\widetilde{k(X)}|k(X)\right)$ is equal to $\mathrm{SL}_2(k)$. We apply Proposition 6.26 in [HS], to deduce that $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)$ is Zariski-dense in $\mathrm{SL}_2(k)$. By Proposition 42 in [C72], we obtain that there are two possibilities:

- $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)=\mathrm{SL}_2(k)$
- $Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)$ is conjugate to $\mathrm{SL}_2(k_0)$ over $\mathrm{SL}_2(k)$, where $k_0=\{a\in k(X)|\partial_Xa=\partial_ta=0\}$.

We see in [D], Remark 4.4, that the last case occurs if and only if the following parameterized differential equation has a solution in $\mathcal{M}_U(X)$, for some polydisc $\varnothing \neq U$ of $\mathbb{C}^n$:

$$\partial_X^3 y(X,t) = \partial_X y(X,t)\frac{4+4tX^3}{X^7} + y(X,t)\partial_X \frac{4+4tX^3}{X^7} - \partial_t \frac{4+4tX^3}{X^7}.$$

With the algorithm presented in [VdPS] p.100, we find that this does not happen and then:

$$Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right) = \mathrm{SL}_2(k).$$

□

Lemma 3.7. *The singular directions of the equation (5) are:*

$$\frac{2k\pi}{5}, \text{with } k \in \mathbb{Z}.$$

Proof. Let $k \in \mathbb{Z}$. The matrix

$$\begin{pmatrix} Xy_k(X^{-1},t) & Xy_{k+1}(X^{-1},t) \\ \partial_X Xy_k(X^{-1},t) & \partial_X Xy_{k+1}(X^{-1},t) \end{pmatrix},$$

is a fundamental solution for the equation $\begin{pmatrix} \partial_X W(X,t) \\ \partial_X^2 W(X,t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \frac{1+tX^3}{X^7} & 0 \end{pmatrix} \begin{pmatrix} W(X,t) \\ \partial_X W(X,t) \end{pmatrix}$. The fundamental solution admits an asymptotic representation on the sectors:

$$\left\{X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]\frac{(2k-1)\pi}{5}, \frac{(2k+3)\pi}{5}\right[\right\}.$$

The only level is $\frac{5}{2}$. From Proposition 1.11 and the construction of the singular directions, we find that the singular directions are

$$\frac{2k\pi}{5}, \text{ with } k \in \mathbb{Z}.$$

□

Example 3.8. We want to compute the Stokes matrix in the direction $\frac{8\pi}{5}$ for the fundamental solution:

$$\begin{pmatrix} Xy_1(X^{-1},t) & Xy_2(X^{-1},t) \\ \partial_X Xy_1(X^{-1},t) & \partial_X Xy_2(X^{-1},t) \end{pmatrix}.$$

We recall the construction of the Stokes matrices. See §1.3 for the notations. Let $\hat{H}(X,t)X^{L(t)}e(Q(X,t))$ be a fundamental solution in the parameterized Hukuhara-Turrittin's canonical form. Let $H^-(X,t)$ (resp. $H^+(X,t)$) be the matrix such that

$$H^-(X,t)e^{L(t)\log(X)}e^{Q(X,t)} \quad \left(\text{resp. } H^+(X,t)e^{L(t)\log(X)}e^{Q(X,t)}\right)$$

is the germ of an asymptotic solution on the sector

$$\left\{X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]\pi, \tfrac{9\pi}{5}\right[\right\} \quad \left(\text{resp. } \left\{X \in \widetilde{\mathbb{C}} \middle| \arg(X) \in \left]\tfrac{7\pi}{5}, \tfrac{11\pi}{5}\right[\right\}\right)$$

The Stokes matrix in the direction $\frac{8\pi}{5}$ is the matrix that sends

$$H^-(X,t)e^{L(t)\log(X)}e^{Q(X,t)} \text{ to } H^+(X,t)e^{L(t)\log(X)}e^{Q(X,t)}.$$

With the domain of definition of the asymptotic representation of $X\widetilde{y}_1(X^{-1},t)$, we deduce from the definition of the Stokes operators that:

$$St_{\frac{8\pi}{5}}(Xy_1(X^{-1},t)) = Xy_1(X^{-1},t). \tag{6}$$

We first write $St_{\frac{8\pi}{5}}(Xy_2(X^{-1},t))$ in the basis

$$(Xy_0(X^{-1},t), Xy_1(X^{-1},t)).$$

There exist $a(t)$ and $b(t) \in \mathcal{M}_U$ such that:

$$St_{\frac{8\pi}{5}}(Xy_2(X^{-1},t)) = a(t)Xy_0(X^{-1},t) + b(t)Xy_1(X^{-1},t).$$

By the construction of the asymptotic solutions with Laplace and Borel transforms (see Proposition 1.9), the asymptotic representation of $St_{\frac{8\pi}{5}}(Xy_2(X^{-1},t))$ has to be bounded in some sector of $\left]\frac{7\pi}{5}, \frac{11\pi}{5}\right[$[2]. Therefore, $a(t) = 0$ or $b(t) = 0$. Since the Stokes operators are automorphisms, we get $b(t) = 0$. Lemma 3.6 says that the parameterized differential Galois group is $\mathrm{SL}_2(k)$. Therefore, because of Proposition 2.20 and Lemma 2.14, the determinant of the matrix has to be 1. Thus by (4), we get that the Stokes matrix in direction $\frac{8\pi}{5}$ is:

$$St_{\frac{8\pi}{5}} = \begin{pmatrix} 1 & -C_0(t)e^{\frac{3i\pi}{5}} \\ 0 & 1 \end{pmatrix}.$$

LEMMA 3.9. *Let $C_0(t)$ be defined as above. Assume that $C_0(t)$ is ∂_t-finite over k. Then, the ∂_t-differential transcendence degree (see §2.1 for definition) of $\widetilde{k(X)}$ over $k(X)$ is at most 2.*

Proof. The extension field $\widetilde{k(X)}$ is generated over $k(X)$ by $y_1(X^{-1},t)$ and $y_2(X^{-1},t)$. By the parameterized Galois correspondence (see Theorem 9.5 in [CS]), the Kolchin closure of the group generated by $St_{\frac{8\pi}{5}}$ is equal to

$$Gal^{\Delta_t}_{\partial_X}\left(\widetilde{k(X)}|F\right),$$

where F is the subfield of $\widetilde{k(X)}$ fixed by $St_{\frac{8\pi}{5}}$. Using (6), we deduce that F contains

$$k(X)\langle y_1(X^{-1},t)\rangle_{\partial_X,\partial_t}.$$

Because $C_0(t)$ satisfies a linear differential equation with coefficients in k, there exists P, a linear differential polynomial such that this group is of the form

$$\left\{\begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix}, \text{ with } P(\alpha) = 0 = P(C_0(t))\right\},$$

and has ∂_t-differential dimension over k equals to 0. Therefore by Proposition 2.7, the ∂_t-differential transcendence degree of $\widetilde{k(X)}$ over F is equal to 0. Because of the fact that F contains $k(X)\langle y_1(X^{-1},t)\rangle_{\partial_X,\partial_t}$, there exists a differential polynomial Q with coefficients in $k(X)$ such that:

$$Q(y_1(X^{-1},t), y_2(X^{-1},t)) = 0 = Q(\partial_X(y_1(X^{-1},t)), \partial_X(y_2(X^{-1},t))).$$

Therefore, the ∂_t-differential transcendence degree of $\widetilde{k(X)}$ over $k(X)$ is at most 2, because $\widetilde{k(X)}$ is generated as a ∂_t-differential field over $k(X)$ by

$$\{y_1(X^{-1},t), y_2(X^{-1},t), \partial_X(y_1(X^{-1},t)), \partial_X(y_2(X^{-1},t))\}.$$

□

THEOREM 3.10. *The function $C_0(t)$ is not ∂_t-finite over k.*

---

[2] This mean that there exist $\frac{7\pi}{5} < \alpha < \beta < \frac{11\pi}{5}$ and $\varepsilon > 0$ such that $St_{\frac{8\pi}{5}}(Xy_2(X^{-1},t))$ is uniformly bounded for $\arg(X) \in]\alpha, \beta[$ and $X < |\varepsilon|$.

Proof. As we see from Lemma 3.6,

$$Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)=\mathrm{SL}_2(k).$$

Therefore, by Proposition 2.7, the ∂_t-differential transcendence degree of $\widetilde{k(X)}$ over $k(X)$ is 3. If $C_0(t)$ is ∂_t-finite over k, because of Lemma 3.9, the ∂_t-differential transcendence degree of $\widetilde{k(X)}$ over $k(X)$ is smaller than 3. Therefore, $C_0(t)$ is not ∂_t-finite over k. ☐

3.3 Linear differential algebraic groups as parameterized differential Galois groups.

As in §3.2, let k be a so-called universal (Δ_t)-field of characteristic 0. Let us consider an equation $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in \mathrm{M}_m(k(X))$, let $\widetilde{k(X)}|k(X)$ be the parameterized Picard-Vessiot extension, and let $G=Gal_{\partial_X}^{\Delta_t}\left(\widetilde{k(X)}|k(X)\right)\subset \mathrm{GL}_m(k)$ be the parameterized differential Galois group defined in §2.1. The following theorem of Seidenberg, applied with $K_0=\mathbb{Q}$ and K_1, the (Δ_t)-field generated by $\mathbb{Q}$ and the X-coefficients of $A(X,t)$, tell us that there exists a polydisc $\varnothing\neq U$, such that $A(X,t)$ may be seen as an element of $\mathrm{M}_m(\mathcal{M}_U(X))$.

Theorem 3.11 (Seidenberg,[Sei58, Sei69]). *Let $\mathbb{Q}\subset K_0\subset K_1$ be finitely generated (Δ_t)-differential extensions of $\mathbb{Q}$, and assume that K_0 consists of meromorphic functions on some domain U of $\mathbb{C}^n$. Then, K_1 is isomorphic to the field K_1^* of meromorphic functions on $U'\subset U$ such that $K_0|_{U'}\subset K_1^*$, and the derivations in Δ_t can be identified with the derivations with respect to the coordinates on U'.*

Let $\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)$ be the parameterized Picard-Vessiot extension defined in §2.5 and let $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ be the parameterized differential Galois group. Using Corollary 2.22, we find that $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$ contains a finitely generated subgroup that is Kolchin-dense in $Aut_{\partial_X}^{\Delta_t}\left(\widetilde{\mathcal{M}_U(X)}|\mathcal{M}_U(X)\right)$. With Proposition 2.20, we find that G contains a finitely generated subgroup that is Kolchin-dense in G. Combined with Corollary 5.2 in [MS11], which gives the sufficiency of the condition, this yields the following result:

Theorem 3.12 (Inverse problem). *Let G be a linear differential algebraic subgroup of $\mathrm{GL}_m(k)$. Then, G is the global parameterized differential Galois group of some equation having coefficients in $k(X)$ if and only if G contains a finitely generated subgroup that is Kolchin-dense in G.*

In the unparameterized case, any linear algebraic group defined over $\mathbb{C}$ is a Galois group of a Picard-Vessiot extension (see [Tre]). In fact, every linear algebraic group defined over $\mathbb{C}$ contains a finitely generated subgroup that is Zariski-dense, which means that Theorem 3.12 is a generalization of the result in [Tre].

The situation is more complicated in the parameterized case. For example, the additive group:

$$\left\{\begin{pmatrix}1 & \alpha\\ 0 & 1\end{pmatrix}, \text{ with } \alpha\in k\right\},$$

is not the global parameterized differential Galois group of any equation having coefficients in $k(X)$ (see Section 7 of [CS]). In the parameterized case with only regular singular poles, the problem has been solved in [MS11], Corollary 5.2: they obtain the same necessary and sufficient condition on the group than Theorem 3.12. In [S11], the author characterizes the linear algebraic subgroups of $\mathrm{GL}_m(k)$ that appears as the global parameterized differential Galois group of some equation having coefficients in $k(X)$: they are the groups such that the identity component has no quotient isomorphic to the additive or multiplicative group of k.

Appendix A.

Let us keep the same notations as in §1.1 and §1.2. The goal of the appendix is to prove the following theorem. See also [BJL, LR01] for proof in the unparameterized case.

Theorem A.1. *Let $\partial_X Y(X,t)=A(X,t)Y(X,t)$, with $A(X,t)\in \mathrm{M}_m(\hat{K}_U)$. There exists a polydisc $\varnothing\neq U'\subset U$, such that we have a fundamental solution of the form*

$$\hat{P}(X,t)X^{C(t)}e(Q(X,t))\in \mathrm{GL}_m(\widehat{\boldsymbol{K}_{U'}}),$$

with:

- $\hat{P}(X,t)\in \mathrm{GL}_m(\hat{K}_{U'})$.
- $C(t)\in \mathrm{M}_m(\mathcal{M}_{U'})$ *and* $X^{C(t)}\in \mathrm{GL}_m(K_{F,U'})$ *satisfy*
$$\partial_X X^{C(t)}=\frac{C(t)}{X}X^{C(t)}=X^{C(t)}\frac{C(t)}{X}.$$
- $e(Q(X,t))=\mathrm{Diag}(e(q_i(X,t)))$, *with* $q_i(X,t)\in \boldsymbol{E}_{U'}$.

Moreover, we may choose the same polydisc $\varnothing\neq U'$ as in Proposition 1.1. Combined with Remark 1.3, this gives a sufficient condition on $t_0\in U$, to have a fundamental solution $\hat{P}(X,t)X^{C(t)}e(Q(X,t))\in \mathrm{GL}_m(\widehat{\boldsymbol{K}_{U'}})$ in the same form as above with $t_0\in U'$.

Notice that contrary to Proposition 1.1, $\hat{H}(X,t)\in \mathrm{GL}_m(\hat{K}_{U'})$. On the other hand, we loose the commutation between $X^{C(t)}$ and $e(Q(X,t))$. Before giving the proof of the theorem, we state and prove two lemmas.

Lemma A.2. *Let $\varnothing\neq U'\subset U$ be a polydisc. Let $a(t)\in\mathcal{M}_{U'}$ and $\alpha(X,t)\in\hat{K}_{F,U'}$ such that $\hat{m}(\alpha(X,t))=a(t)\alpha(X,t)$. Then there exists $\hat{h}(X,t)\in\hat{K}_{U'}$ and $b(t)\in\mathcal{M}_{U'}$ such that $\alpha(X,t)=\hat{h}(X,t)X^{b(t)}$.*

Proof. Let $\alpha(X,t)\in\hat{K}_{F,U'}$ such that $\hat{m}(\alpha(X,t))=a(t)\alpha(X,t)$. The element $\alpha(X,t)$ belongs to the fraction field of a free polynomial ring:

$$P=\hat{K}_{U'}[\log, X^{b_1(t)},\dots,X^{b_k(t)}].$$

Write $\alpha(X,t)=\alpha_1(X,t)/\alpha_2(X,t)$ with gcd in P equals to 1. Using the relations in $\hat{K}_{F,U'}$, and applying $\hat{m}$ to $\alpha_1(X,t)/\alpha_2(X,t)$, we find that $\alpha(X,t)$ contains no terms in log. One can normalize $\alpha_2(X,t)$ such that it contains a term of the form $X^{n_1b_1(t)+\dots+n_kb_k(t)}$ with coefficient 1 and $n_i\in\mathbb{Z}$. Using $\hat{m}(\alpha_1(X,t)/\alpha_2(X,t))=a(t)\alpha_1(X,t)/\alpha_2(X,t)$, we find that $\hat{m}(\alpha_2(X,t))=e^{2i\pi(n_1b_1(t)+\dots+n_kb_k(t))}\alpha_2(X,t)$ and $\hat{m}(\alpha_1(X,t))=a(t)e^{2i\pi(n_1b_1(t)+\dots+n_1b_1(t))}\alpha_1(X,t)$, which is impossible, unless

$$e^{2i\pi(n_1b_1(t)+\dots+n_kb_k(t))}=1.$$

This means that $\alpha_2(X,t)\in\hat{K}_{U'}$ and we may assume $\alpha_2(X,t)=1$. Applying $\hat{m}$ to $\alpha_1(X,t)$, one finds that $\alpha_1(X,t)$ contains at most one term, that is $\alpha(X,t)=\hat{h}(X,t)X^{b(t)}$, with $\hat{h}(X,t)\in\hat{K}_{U'}$ and $b(t)\in\mathcal{M}_{U'}$ that satisfies $e^{2i\pi b(t)}=a(t)$. □

Lemma A.3. *Let* $\varnothing \neq U' \subset U$ *be a polydisc. Let* $A(X,t) \in \mathrm{M}_m(\hat{K}_{U'})$. *Let* $F_1(X,t)e(Q_1(X,t))$ *and* $F_2(X,t)e(Q_2(X,t))$ *be two fundamental solutions of:*

$$\partial_X Y(X,t) = A(X,t)Y(X,t),$$

satisfying, for $i \in \{1;2\}$, $F_i(X,t) \in \mathrm{GL}_m(\hat{K}_{F,U'})$ *and* $Q_i(X,t) = \mathrm{Diag}[q_{i,j}(X,t)]$ *such that* $q_{i,j}(X,t)$ *belongs to* $\boldsymbol{E}_{U'}$. *Then,* $F_1(X,t)^{-1}F_2(X,t) \in \mathrm{GL}_m(\mathcal{M}_{U'})$.

Proof. A straightforward computation shows that:

$$\partial_X \left(F_1(X,t)e(Q_1(X,t))\right)^{-1}\left(F_2(X,t)e(Q_2(X,t))\right) = 0.$$

By Proposition 2.16,

$$\left(F_1(X,t)e(Q_1(X,t))\right)^{-1}\left(F_2(X,t)e(Q_2(X,t))\right) = C(t) \in \mathrm{GL}_m(\mathcal{M}_{U'}).$$

We have the equality:

$$e(Q_1(X,t))C(t)e(-Q_2(X,t)) = F_1(X,t)^{-1}F_2(X,t).$$

The entries of $e(Q_1(X,t))C(t)e(-Q_2(X,t))$ are of the form $C_{i,j}(t)e(q_{1,j}(X,t) - q_{2,j}(X,t))$, with $C_{i,j}(t) \in \mathcal{M}_{U'}$, and the matrix $F_1(X,t)^{-1}F_2(X,t)$ belongs to $\mathrm{GL}_m(\hat{K}_{F,U'})$. By construction,

$$\hat{K}_{F,U'} \cap \mathcal{M}_{U'}((e(q(X,t)))_{q(X,t)\in \mathbf{E}_{U'}}) = \mathcal{M}_{U'},$$

and we obtain that:

$$F_1(X,t)^{-1}F_2(X,t) \in \mathrm{GL}_m(\mathcal{M}_{U'}).$$

□

Proof of Theorem A.1. By Proposition 1.1, we know that we have a fundamental solution of the parameterized linear differential equation $\partial_X Y(X,t) = A(X,t)Y(X,t)$ of the form:

$$\hat{H}(X,t)X^{L(t)}e(Q(X,t)),$$

with $\hat{H}(X,t) \in \mathrm{GL}_m(\hat{K}_{U'}[X^{1/\nu}])$ and $\nu \in \mathbb{N}^*$. From Definition 2.11, $\hat{m}$ commutes with the derivation ∂_X, and therefore $\hat{m}\left(\hat{H}(X,t)X^{L(t)}e(Q(X,t))\right)$ is another fundamental solution. From the construction of $\hat{m}$, we deduce that $\hat{m}\left(\hat{H}(X,t)X^{L(t)}\right) \in \mathrm{GL}_m(\hat{K}_{F,U'})$, and we can apply Lemma A.3 to deduce the existence of $\hat{M}(t) \in \mathrm{GL}_m(\mathcal{M}_{U'})$ such that:

$$\hat{m}\left(\hat{H}(X,t)X^{L(t)}\right) = \hat{H}(X,t)X^{L(t)}\hat{M}(t). \tag{7}$$

Let $\hat{M}(t) = D(t)U(t)$, with $D(t)$ diagonalizable and $U(t)$ unipotent such that $D(t)U(t) = U(t)D(t)$ is the multiplicative analogue of the Jordan decomposition of $\hat{M}(t)$. If $a(t)$ is an eigenvalue of $D(t)$ (and therefore $a(t)$ is an eigenvalue of $\hat{M}(t)$), then there exists $0 \neq \alpha(X,t) \in \hat{K}_{F,U'}$ such that $\hat{m}(\alpha(X,t)) = a(t)\alpha(X,t)$, because of the relation (7). By Lemma A.2, $\alpha(X,t)$ is equal to $\hat{h}(X,t)X^{b(t)}$, with $b(t) \in \mathcal{M}_{U'}$ satisfying $e^{2i\pi b(t)} = a(t)$ and $\hat{h}(X,t) \in \hat{K}_{U'}$. This implies that $a(t)$ and all the eigenvalues of $D(t)$ are of the form $e^{\beta(t)}$, with $\beta(t) \in \mathcal{M}_{U'}$. So we have proved the existence of $C(t) \in \mathrm{M}_m(\mathcal{M}_{U'})$ such that $e^{2i\pi C(t)} = \hat{M}(t)$. Let:

$$\hat{P}(X,t) = \hat{H}(X,t)X^{L(t)}X^{-C(t)}.$$

A computation shows that the monodromy of $X^{C(t)}$ is:

$$\hat{m}\left(X^{C(t)}\right) = e^{2i\pi C(t)}X^{C(t)} = X^{C(t)}e^{2i\pi C(t)}.$$

The matrix $\hat{P}(X,t)$ is fixed by the monodromy and therefore belongs to $\mathrm{GL}_m(\hat{K}_{U'})$, because of Proposition 2.16. Finally,

$$\hat{P}(X,t)X^{C(t)}e(Q(X,t))$$

is a fundamental solution of the parameterized linear differential equation $\partial_X Y(X,t) = A(X,t)Y(X,t)$ in the wished form. $\square$

References

A Acosta-Humanez, Primitivo B, Galoisian Approach to Supersymmetric Quantum Mechanics, Phd Dissertation, Universitat Politecnica de Catalunya, 2009.

Ar Carlos E. Arreche , Computing the differential Galois group of a one-parameter family of second order linear differential equations.

B Balser, Werner, From divergent power series to analytic functions, Lecture Notes in Mathematics, 1582, Theory and application of multisummable power series, Springer-Verlag, Berlin, 1994, 3-540-58268-1.

Ban S.B. Bank, Some results on hypertranscendental meromorphic functions. Monatsh. Math. 90 (1980), no. 4, 267-289.

Ber Bertrand, Daniel, Groupes algébriques et équations différentielles linéaires, Séminaire Bourbaki, Vol. 1991/92, Astérisque, Astérisque, 206, 1992, Exp. No. 750, 4, 183–204.

BJL Balser, W. and Jurkat, W. B. and Lutz, D. A., A general theory of invariants for meromorphic differential equations. I. Formal invariants, Funkcial. Ekvac., Fako de l'Funkcialaj Ekvacioj Japana Matematika Societo. Funkcialaj Ekvaciog. Serio Internacia, 22, 1979, 2, 197–221.

Bol A.A. Bolibruch, On isomonodromic deformations of Fuchsian systems. J. Dynam. Control Systems 3 (1997), no. 4, 589-604.

BV Babbitt, Donald G. and Varadarajan, V. S., Deformations of nilpotent matrices over rings and reduction of analytic families of meromorphic differential equations, Mem. Amer. Math. Soc., 55, 1985, 325, iv+147.

C72 P.J. Cassidy, Differential algebraic groups. Amer. J. Math. 94 (1972), 891-954.

C89 P.J. Cassidy, The classification of the semisimple differential algebraic groups and the linear semisimple differential algebraic Lie algebras. J. Algebra 121 (1989), no. 1, 169-238.

CHS Z. Chatzidakis/ C. Hardouin/ M.F. Singer, On the definitions of difference Galois groups. Model theory with applications to algebra and analysis. Vol. 1, 73-109, London Math. Soc. Lecture Note Ser., 349, Cambridge Univ. Press, Cambridge, 2008.

CR J. Cano / J.P. Ramis. Théorie de Galois différentielle, multisommabilité et phénomène de Stokes. Preprint.

CS Cassidy, Phyllis J./Singer, Michael F., Galois theory of parameterized differential equations and linear. differential algebraic groups, Differential equations and quantum groups, IRMA Lect. Math. Theor. Phys., 9, 2007, 113-155.

DVH Di Vizio, Lucia and Hardouin, Charlotte, Descent for differential Galois theory of difference equations: confluence and q-dependence, Pacific J. Math., Pacific Journal of Mathematics, 256, 2012, 1, 79–104.

D T. Dreyfus, The Kovacic's algorithm for parameterized differential Galois theory, preprint. URL http://arxiv.org/pdf/1110.1053.pdf .

E J. Ecalle. Résurgence et accélération. Cours de 3 ème cycle.

E81 J.Écalle, Les algébres de fonctions résurgentes, Publ. Math. D'Orsay, Université Paris-Sud, 1981.05 (1981).

GGO H.Gillet/S. Gorchinskiy/A.Ovchinnikov, Parameterized Picard Vessiot extensions and Atiyah extensions, Preprint, october 2011. URL http://arxiv.org/abs/1110.3526 .

GO S. Gorchinskiy/A.Ovchinnikov, Isomonodromic differential equations and differential Tannakian categories, Preprint, 2012. URL http://arxiv.org/abs/1202.0927

HS C. Hardouin/ M.F. Singer, Differential Galois theory of linear difference equations.Math. Ann. 342 (2008), no. 2, 333-377.

Kol76 E.R. Kolchin, Differential algebra and Algebraic groups, Academic Press, New York 1976.

Kol85 E.R. Kolchin, Differential algebraic groups. Pure and Applied Mathematics, 114. Academic Press, Inc., Orlando, FL, 1985. xvii+271 pp. ISBN: 0-12-417640-2.

Kov J.J.Kovacic, An algorithm for solving second order linear homogeneous differential equations. J. Symbolic Comput. 2 (1986), no. 1, 3-43.

L Landesman, Peter, Generalized differential Galois theory, Trans. Amer. Math. Soc., Transactions of the American Mathematical Society, 360, 2008, 8, 4441–4495, 0002-9947.

LR90 M Loday-Richaud, Introductiona la multisommabilité, Gazette des Mathématiciens, SMF, 1990.

LR94 Loday-Richaud, Michèle, Stokes phenomenon, multisummability and differential Galois groups, Ann. Inst. Fourier (Grenoble), Université de Grenoble. Annales de l'Institut Fourier, 44, 1994, 3, 849–906.

LR95 M.Loday-Richaud, Michèle; Solutions formelles des systèmes différentiels linéaires méromorphes et sommation. Exposition. Math. 13 (1995), no. 2/3, 116-162.

LR01 Loday-Richaud, Michèle, Rank reduction, normal forms and Stokes matrices, Expo. Math., Expositiones Mathematicae, 19, 2001, 3, 229–250.

LRR Loday-Richaud, Michèle and Remy, Pascal, Resurgence, Stokes phenomenon and alien derivatives for level-one linear differential systems, J. Differential Equations, Journal of Differential Equations, 250, 2011, 3, 1591–1630, 0022-0396, http://dx.doi.org/10.1016/j.jde.2010.09.018.

LT Laurent-Thiébaut, Christine, Théorie des fonctions holomorphes de plusieurs variables, French, Savoirs Actuels. [Current Scholarship], Mathématiques. [Mathematics], InterEditions, Paris, 1997,xiv+245, 2-7296-0660-2.

Ma Magid, Andy R., Lectures on differential Galois theory, University Lecture Series, 7, American Mathematical Society, Providence, RI, 1994, xiv+105, 0-8218-7004-1.

M82 B.Malgrange, Sur les déformations isomonodromiques. Singularités régulières. Mathematics and physics (Paris, 1979/1982), 401-426, Progr. Math., 37, Birkhäuser Boston, Boston, MA, 1983.

M91 B. Malgrange, Équations différentielles à coefficients polynomiaux. Progress in Mathematics, 96. Birkhäuser Boston, Inc., Boston, MA, 1991. vi+232 pp. ISBN: 0-8176-3556-4.

M95 B. Malgrange, Sommation des séries divergentes. Exposition. Math. 13 (1995), no. 2-3, 163-222.

M02 B.Malgrange, On nonlinear differential Galois theory. Dedicated to the memory of Jacques-Louis Lions. Chinese Ann. Math. Ser. B 23 (2002), no. 2, 219-226.

MO A. Minchenko and A. Ovchinnikov, Zariski Closures of Linear Differential Algebraic Groups, to appear in Advances in Mathematics, 10.1016/j.aim.2011.03.002.

MR B.Malgrange/J. Ramis, Fonctions multisommables. Ann. Inst. Fourier (Grenoble) 42 (1992), no. 1-2, 353-368.

M Mitschi, Claude, Differential Galois groups of confluent generalized hypergeometric equations: an approach using Stokes multipliers, Pacific J. Math., 176, 1996, 2, 365–405.

MS C.Mitschi/ M.F. Singer, Projective Isomonodromy and Galois Groups: to appear in Proceedings of the Amer. Mat. Soc.

MS11 Monodromy groups of parameterized linear differential equations with regular singularities Bull. London Math. Soc. (2012) 44(5).

N Peón Nieto, Ana, On $\sigma\delta$-Picard-Vessiot extensions, Comm. Algebra, 39, 2011, 4, 1242–1249.

R79 J.P. Ramis, Les séries k-sommables et leurs applications, Analysis, Microlocal Calculus and Relativistic Quantum Theory, Proceeding Les Houches (1979), Springer Notes in Physics 126, p. 178-199 (1980).

R85 J.P Ramis, Phénomène de Stokes et filtration Gevrey sur le groupe de Picard-Vessiot. C. R. Acad. Sci. Paris Sér. I Math. 301 (1985), no. 5, 165-167.

RM1 Ramis, J.-P / Martinet, J., Théorie de Galois différentielle et resommation, Computer algebra and differential equations, Comput. Math. Appl., 117–214, Academic Press, London, 1990.

Ras J-M Rasoamanana, Résurgence-sommabilité de séries formelles ramifiées dépendant d'un paramètre et solutions d'équations différentielles linéaires. Ann. Fac. Sci. Toulouse Math. (6) 19 (2010), no. 2, 303-343, 34/XX.

Re Remy, Pascal, Matrices de Stokes-Ramis et constantes de connexion pour les systèmes différentiels linéaires de niveau unique, Ann. Fac. Sc. Toulouse, XXI-2 (2012), 93-150.

Rob A. Robinson, On the concept of a differentially closed field. Bull. Res. Council Israel Sect. F 8F 1959 113/128 (1959).

RS J.P. Ramis / Y. Sibuya. Hukuhara's domains and fundamental existence and uniqueness theorems for asymptotics solutions of Gevrey type, Asympt. Anal. 2 (1989), 39-94.

S09 Singer, Michael F.,Introduction to the Galois theory of linear differential equations, Algebraic theory of differential equations, London Math. Soc. Lecture Note Ser., 357,Cambridge Univ. Press,Cambridge,2009,1–82.

S11 M.F. Singer, Linear algebraic groups as parameterized Picard-Vessiot Galois groups. Preprint. URL http://www4.ncsu.edu/∼singer/papers/Singer_LinGpsPPV.pdf .

Sch Schäfke, Reinhard Formal fundamental solutions of irregular singular differential equations depending upon parameters. J. Dynam. Control Systems 7 (2001), no. 4, 501-533.

Sei58 Seidenberg, A., Abstract differential algebra and the analytic case, Proc. Amer. Math. Soc., Proceedings of the American Mathematical Society, 9, 1958, 159–164.

Sei69 Seidenberg, A., Abstract differential algebra and the analytic case. II, Proc. Amer. Math. Soc., Proceedings of the American Mathematical Society, 23, 1969, 689–691.

Si Y. Sibuya, Global theory of a second order linear ordinary differential equation with a polynomial coefficient. North-Holland Mathematics Studies, Vol. 18. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975. xv+290 pp. ISBN: 0-7204-2609-X.

Si90 Y. Sibuya, Linear differential equations in the complex domain: problems of analytic continuation. Translated from the Japanese by the author. Translations of Mathematical Monographs, 82. American Mathematical Society, Providence, RI, 1990. xiv+269 pp. ISBN: 0-8218-4535-7.

Tre Tretkoff, Carol and Tretkoff, Marvin, Solution of the inverse problem of differential Galois theory in the classical case, Amer. J. Math., American Journal of Mathematics, 101, 1979, 6, 1327–1332.

Um H.Umemura, Galois theory of algebraic and differential equations. Nagoya Math. J. 144 (1996), 1-58.

VdPS van der Put, M/ Singer, Michael F. , Galois theory of linear differential equations, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 328, Springer-Verlag, Berlin, 2003, 3-540-44228-6.

W Wasow, Wolfgang, Asymptotic expansions for ordinary differential equations, Dover Publications Inc., New York, 1987, x+374, 0-486-65456-7.

Wat Watson, G. N., A treatise on the theory of Bessel functions, Cambridge University Press, Cambridge, England; The Macmillan Company, New York, 1944. vi+804 pp.

Wib M.Wibmer, Existence of ∂-parameterized Picard-Vessiot extensions over fields with algebraically closed constants, arXiv:1104.3514.

Thomas Dreyfus `tdreyfus@math.jussieu.fr.`
Institut de mathématiques de Jussieu, Topologie et géométrie algébriques
Current address: 4 place Jussieu 75005 Paris.